

FORM PTO-1390  
(REV 12-29-99)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

4512/00012

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/744502

INTERNATIONAL APPLICATION NO.

PCT/GB99/02406

INTERNATIONAL FILING DATE

23 July 1999 (23.07.99)

PRIORITY DATE CLAIMED

24 July 1998 (24.07.98)

TITLE OF INVENTION Protection of Plants using Rhizoctonia

APPLICANT(S) FOR DO/EO/US RUBIO SUSAN, Victor; SALAZAR TORRES, Oscar; JULIAN ESQUIVIAS, Maria; GONZALES GARCIA, Vicente;  
GOMEZ-ACEBA GULLON, Eduardo; MUNOZ GOMEZ, Ramona; LOPEZ CORCOLES, Horacio

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

## Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.  
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

Also enclosed are copies of the following documents:

- . PCT Request Form
- . International Search Report
- . International Application (PCT Publication)
- . PCT Demand
- . International Preliminary Examination Report
- . Deposit Receipts (5 Sheets)

09/744502

PCT/GB99/02406

4512/00012

17. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):**

Neither international preliminary examination fee (37 CFR 1.482)  
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO  
and International Search Report not prepared by the EPO or JPO ..... \$970.00

International preliminary examination fee (37 CFR 1.482) not paid to  
USPTO but International Search Report prepared by the EPO or JPO ..... \$840.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but  
international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$670.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$96.00

**ENTER APPROPRIATE BASIC FEE AMOUNT =****CALCULATIONS** PTO USE ONLY

\$ 860.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$ 0.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	10 - 20 =	0	X \$18.00
Independent claims	3 - 3 =	0	X \$78.00

\$ 0.00

\$ 0.00

MULTIPLE DEPENDENT CLAIM(S) (if applicable)

+ \$260.00

\$ 0.00

**TOTAL OF ABOVE CALCULATIONS =**

\$ 860.00

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement  
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

\$ 0.00

**SUBTOTAL =**

\$ 860.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$ 0.00

**TOTAL NATIONAL FEE =**

\$ 860.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be  
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

\$ 0.00

**TOTAL FEES ENCLOSED =**

\$ 860.00

Amount to be  
refunded:

\$

charged:

\$ 860.00

a. ☐ A check in the amount of \$\_\_\_\_\_ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 19-0733 in the amount of \$ 860.00 to cover the above fees.  
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
overpayment to Deposit Account No. 19-0733. A duplicate copy of this sheet is enclosed.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Ernest V. Linek  
BANNER & WITCOFF, LTD.  
28 State Street, 28th Floor  
Boston, MA 02109

SIGNATURE:

Peter D. McDermott

NAME

29,411

REGISTRATION NUMBER

64/ PPT  
500 Rec'd PCT/PTO 24 JAN 2001PROTECTION OF PLANTS USING RHIZOCTONIA

The present invention describes a new group of binucleate *Rhizoctonia* suited for use in plant protection.

#### Background Of The Invention

The complex of species *Rhizoctonia* represent ubiquitous and variable fungi that inhabit soil. Presently there are about 120 species or varieties. A major taxonomic feature is the presence of either multinucleate or binucleate cells in young vegetative hyphae. The multinucleate *Rhizoctonia solani* presents the teleomorph *Thanatephorus cucumeris*, while the binucleate *Rhizoctonia* species include strains presenting a *Ceratobasidium* species teleomorph. Many different types of plant pathogenic *Rhizoctonia* cause economically significant diseases on most of the important crops worldwide.

Thus, *Rhizoctonia* diseases in field, vegetable, ornamental, nursery and greenhouse crops cause large losses to growers, which result in increased costs to consumers. Strains of *Rhizoctonia solani* Kühn, and certain other *Rhizoctonia* species are capable of attacking a wide range of hosts, causing seed decay, damping-off of seedlings, stem cankers, root rots, fruit decays and foliage diseases. Moreover, *Rhizoctonia solani* occurs worldwide in most cultivated soils and is indigenous in many uncultivated areas.

At present, *Rhizoctonia* diseases on some hosts are only treatable with fungicides. For small acreages of high valuable crops, such as greenhouse and nursery crops, pathogen eradication by heat treatment or chemical fumigation is useful, but re-infestation often presents renewed problems prior to harvest. Fungicides pose risk to health, water quality and other aspects of the environment. Additionally some fungicides vary in effectiveness among and within anastomosis groups (AG) of *Rhizoctonia solani*, suggesting that their extensive use could result in development of resistance as occurs within other plant pathogen populations.

Despite research on biological control dating back to the early 1930s, reliable, economical biocontrol of diseases caused by *Rhizoctonia solani* are not generally available on a commercial scale. New approaches for detection and use of novel agents and development of broadly applicable biological control management systems are needed, specially for field crops. During the past 10 years, new sources of agents from within the diverse groups of binucleate *Rhizoctonia* spp. and hypovirulent *Rhizoctonia solani* isolates have been demonstrated to be effective in biocontrol of a range of host - *Rhizoctonia solani* disease combinations. Tested collections of binucleate *Rhizoctonia* spp. and hypovirulent *Rhizoctonia solani* differ markedly in effectiveness as biocontrol agents for diseases caused by *Rhizoctonia solani*.

In the book *Rhizoctonia* Species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control, edited by Sneh, Jabaji-Hare and Dijst, published by Kluwer Academic Publishers, Dordrecht in 1996, there is a Chapter VI.B.4 by B. Sneh on Nonpathogenic Isolates of *Rhizoctonia* (np-R) spp. and Their Role in Biological Control, with a survey of the literature at that time. A subsequent review by B. Sneh is in Biotechnology Advances, 16: 1-32 (1998), on Use of Non-pathogenic or Hypovirulent Fungal Strains to Protect Plants against Closely Related Fungal Pathogens.

The use of non-pathogenic *Rhizoctonia* strains as biocontrol agents of crop plant diseases has a great potential to increase crop yields but there are some problems in identifying candidate strains. Indeed, more generally the classification of *Rhizoctonia* is replete with difficulties.

Various methods have been employed to classify isolates of *Rhizoctonia*, relying for example on their ecology, enzymology, host range specificity, morphology and hyphal anastomosis. The morphological characteristics of anamorph and teleomorph of binucleate species are very similar, and reliable identification is often difficult.

The identification at strain level of *Rhizoctonia solani* AG4 isolates by direct sequence of asymmetric products of the ITS regions is reported in Curr Genet (1996); 29: 174 to 181. The relatedness of nine isolates of *Rhizoctonia solani*, belonging to anastomosis group (AG) 4, and



one isolate of AG1, was determined by comparative sequence analysis based on direct sequencing of PCR-amplified ribosomal DNA [the internal transcribed spacer (ITS) region and the 5.8s ribosomal DNA]. The 5.8s ribosomal DNA was completely conserved, but both ITS regions showed variation among strains. AG1 was an outgroup based on anastomosis ability and RFLP analyses. Phylogenetic analyses based on the ITS sequences suggested that the analyzed AG4 strains can be divided into three groups that correlated with habitat and virulence.

Sequence variation of the rDNA ITS regions within and between anastomosis groups in *Rhizoctonia solani* is described in Curr Genet (1997) 32: 237 to 243. The authors report that sequence analysis of the rDNA region containing the internal transcribed spacer (ITS) regions and the 5.8s rDNA coding sequence was used to evaluate the genetic diversity of 45 isolates. They report that their results suggest that such sequence analysis may be a valuable tool for identifying AG subgroups of biological significance.

In *Rhizoctonia solani* AG3, the ITS sequences are almost identical in more than 30 isolates we have studied. Therefore such ITS sequence are not suitable for identification in this situation.

Until now there was only one molecular work dealing with binucleate *Rhizoctonia*, the work described by Cubeta *et al.* In this work, in which restriction analysis of the amplified 25 S ribosomal gene was performed to differentiate among anastomosis groups of binucleate *Rhizoctonia*, it was not possible to differentiate between many of the AGs. For instance, AG C, AG F, AG H, AG I and AG J are grouped together and cannot be differentiated, and the same is true for AG A, AG Bo, AG L, AG O, AG K and AG E. When this methodology of rDNA-RFLP was applied to *R. solani*, a considerable molecular variation among and within anastomosis and intraspecific groups was found.

There is a need for a procedure to select isolates or cultures of *Rhizoctonia* which can protect plants against pathogenic fungi.

## Summary Of The Invention

The present invention provides for the protection of plants against pathogenic fungi by the use of binucleate *Rhizoctonia* selected through the molecular detection of certain ITS ribosomal sequences, the protective strains being members of a phylogenetically differentiated cluster.

Thus, the present invention describes a new group of binucleate *Rhizoctonia* able to protect different kind of plants, like tomatoes, wheat, rape, alfalfa, sugar beet, carrot, garlic, onions, grass, pine trees, and others, from infections produced by different pathogenic fungi among them *Fusarium*, *Penicillium* and pathogenic *Rhizoctonia*, and the appropriated methodology for the molecular identification of the new group, its detection based on probes specially designed and its practical application for the protection of crops. More generally, the present invention is of use for protection in a field crop, water bed crop, container-grown crop, vegetable or fruit crop, ornamental crop, nursery crop, garden crop, forest crop, or greenhouse crop.

Thus the present invention uses the ITS data, especially the ITS1 data, to distinguish isolates and in particular to identify pathogenic and protective isolates for each plant belonging to the binucleate group.

The invention also provides specific PCR primers for the ITS molecular analysis of pathogenic and protective strains of *Rhizoctonia*, and which serve to characterise *Rhizoctonia* strains in accordance with this invention.

## Preferred Embodiments

In the present invention, the ITS sequence is employed as a marker to distinguish strains and correspondingly classify the pathogenicity of isolates of *Rhizoctonia*. The ITS information enables phylogenetic location of protective isolates, and design of specific primers for the identification of new isolates. In particular, preferred strains of this invention incorporate

include those obtained by PCR using at least one of the following sequences SEQ ID NO: 1 to SEQ ID NO: 14.

Particularly preferred strains include those with homology to one or more of SEQ ID NO: 7 to SEQ ID NO: 14. Strict homology with these sequences is not ordinarily necessary, and variation by substitution, insertion and/or deletion of say 1, 2 or 3 bases is permitted.

Suitable isolated strains of this invention include the deposited strains *Rhizoctonia* sp. CECT 20320, *Rhizoctonia* sp. CECT 20321, *Rhizoctonia* sp. CECT 20322, *Rhizoctonia* sp. CECT 20323, and *Rhizoctonia* sp. CECT 20324. Other suitable strains include strains which cluster with these deposited strains, when using the methodology herein. More generally, preferred isolates are those which cluster with the isolates of this invention within the clusters shown in Figure 2.

Isolated strains of this invention obtained by us have the ITS1 sequences included in Figure 1, as follows: ID SEQ:NO 15 to ID SEQ:ID 28.

The present invention extends to any strains which have such an ITS1 sequence or at least 90% homology with such a sequence. At least 95% homology is preferred. Moreover, the invention includes strains which include fragments of any one of the sequences SEQ ID NO: 15 to 28. Such fragments may comprise at least 5, 10, 15 or 25 or more nucleic acids, and may be fragments having at least 90%, more preferably at least 95 or 100% homology. More generally, the present invention employs such fragments as signature sequences to identify the strains of the present invention. By their nature, signature sequences serve to uniquely identify the strains of this invention. Further preferred strains include those which have such an ITS2 sequence similar to that in the isolates of this invention, or at least 90% homology with such a sequence. At least 95% homology is preferred.

The protective strain can be utilised in a form of powder obtained after fermentation of the micro-organisms with subsequent drying of the filtered broth, applied to the seed before seeding or applied to the soil before seeding or both, and will preclude the growth of the pathogen *Rhizoctonia* protecting the plant from the infection. Thus, for example, harvested

mycelium from a protective isolate in accordance with this invention can be applied direct to seeds, bulbs or other germinatable form, or applied to the soil or other growth medium, pre- or post-emergence. Suitable dosing levels may be found by experiment, and will vary depending on the mode of administration and the crop to be protected, among other factors. Guidance is to be found in the examples of this invention when taken with the accompanying figures. The present invention further provides concentrates of the mycelium ready for admixing with diluent to form a mix suited for application. The concentrates can be liquid or solid, and suitably contain up to 75% mycelium, typically 10 to 50% mycelium. The strains of this invention are primarily intended for large scale agricultural use, but may also be employed by smaller enterprises and by the public.

Examples of crops which might be protected by the present invention include tomato, pepper, cucumber, lettuce, radish, parsley and other herbs, bean, potato, sugar beet, carrot, garlic, onions, grass, wheat, rape, alfalfa, pine and other trees, as well as flowers.

#### Summary Of Figures

Figure 1. Sequence data for isolates of this invention, aligned with other isolates for comparison. The sequence extends beyond the end of the ITS1 region to the 5.8s region.

Figure 2. Phylogenetic tree generated from a branch and bound algorithm in PAUP 3.1.1, using CLUSTALW for alignment. The tree was rooted with the ITS sequence of a *Trichoderma harzianum* isolate. The numbers above internodes indicate the percentage corresponding to the frequency for which a given branch appeared in a 1000 bootstrap replicates. Horizontal lengths represent genetic distances.

Figures 3 to 10. The results for experiments in the protection of grass, tomato, carrot, wheat, rape, alfalfa, sugar beet and onion are plotted as histograms. The protective effect is expressed as a coefficient on the Butler scale.

Figures 11 to 13. The results for the wheat, alfalfa and sugar beet tests, plotted as histograms.

Figures 14 and 15. Protection test 1.

Figures 16 to 24. Protection test 3.

Figure 25 to 31. Protection test 4.

Figure 32 to 34. Protection test 5.

Figures 35 to 39. Protection test 6.

Figure 40 to 42. Protection test 7.

Figure 43 to 48. Protection test 8.

Figure 49 to 53. Protection test 9.

#### Examples Of The Invention

We sequenced the ITS - 5.8S regions of 13 binucleate isolates of *Rhizoctonia* isolated from Albacete (South-East Spain), 9 binucleate isolates from CBS (Central Bureau Schimmelcultures, Baarn, Delft, Netherlands) collection, 3 multinucleate isolates from Albacete and multinucleate isolates belonging to AG 2. The later multinucleate isolates were included to determine the sequence variation in rDNA regions and to assess this variation as a possible tool for identifying groups into *Rhizoctonia* spp.

Samples of saffron, alfalfa and pine seedlings were collected at different locations in Albacete, South East of Spain. Isolation of *Rhizoctonia* spp. from these samples was performed by washing saffron roots, alfalfa stems and roots and whole pine seedlings with tap water to remove adhering soils particles and plating plant segments on 2% water agar amended with chloramphenicol (250 mg/ml). The cultures were characterised as *Rhizoctonia* by observing

hyphal and colony morphology and vegetative features microscopically, such as hyphae with dolipore septa, basally constricted branching near the distal septum of cells, and absence of clamp connections and conidia. Cultures were maintained and stored at room temperature on potato-dextrose-agar [PDA, Oxoid LTD, U.K.].

The number of nuclei per hyphal cells was determined using a staining method described by Julian *et al.* Briefly, 3 mm<sup>2</sup> agar plugs of inoculum were placed in petri dishes containing liquid complete medium [CM: 5 g malt extract (Merck, Darmstadt, Germany), 5 g yeast extract (Difco, Detroit, USA) and 5 g glucose anhydrous (Merck, Darmstadt, Germany) per liter] and incubated for 3 d at 23°C in the dark. The resulting mycelial colony was transferred to a microscope slide, excess medium was removed and a drop of 50 mg/ml Hoechst 33342 (Molecular Probes Europe, Leiden, The Netherlands) was added. After a 15 min incubation in the dark, the staining solution was removed and 50% glycerol was added. A coverslip was placed on sample and the number of nuclei was determined by fluorescence microscopy using an Olympus BHS microscope supplemented with a BH2-RFC reflected light fluorescence attachment (with dichroic mirrors BH2-DMU for UV excitation).

The isolates obtained are listed in the Table 1. All the isolates have been identified as belonging to the *Ceratobasidium* spp. complex. Other isolates included in this study are binucleate *Rhizoctonia* AG testers. All isolates have been described to present a teleomorph in the genus *Ceratobasidium*, except for AG 1 and AG N, for which a teleomorph is not known. Isolate 521 was obtained from ATCC (code ATCC 64643) and had previously been described as hypovirulent AG 4 and isolate Rh2815 was obtained from Colección Española de Cultivos Tipo (code CECT 2815) and was also described as AG 4.

Binucleate *Rhizoctonia* spp. isolates have been obtained from the CBS collection and are listed in Table 2.

Table 1. Original isolates described and sequenced for the first time in the invention as protective BNR, nuclear condition, host, size of ITS1, ITS2 and 5.8s rDNA gene, and EMBL accession number.

Isolate	Nuclear condition	Host	ITS 1	ITS2	Total	Accession No.
Eab-F1	Binucleate	Saffron	245	275	675	AJ242875
Eab-F2	Binucleate	Saffron	242	260	657	AJ242876
Eab-F3	Binucleate	Saffron	243	260	658	AJ242877
Eab-FS	Binucleate	Saffron	238	261	654	AJ242878
Eab-F6	Binucleate	Saffron	246	251	652	AJ242879
Eab-F7	Binucleate	Saffron	252	254	661	AJ242880
Eab-SI	Binucleate	Saffron	230	280	665	AJ242881
Eab-S2	Binucleate	Saffron	232	254	641	AJ242882
Eab-S3	Binucleate	Saffron	232	260	647	AJ242883
Eab-S4	Binucleate	Saffron	226	254	635	AJ242884
Eab-S5	Binucleate	Saffron	236	269	660	AJ242885
Eab-S6	Binucleate	Pine	237	269	661	AJ242886
Eab-S7	Multinucleate	Pine	241	273	669	AJ242887
Eab-A1	Multinucleate	Alfalfa	274	298	727	AJ242888
Eab-A2	Multinucleate	Alfalfa	273	294	722	AJ242889

Table 2 Reference isolates belonging to the Binucleate Group obtained from different collections

Isolates	Nuclear Condition	Identification	Isolated in	Geographical Origin
521	Binucleate	Unknown	Soil	Israel
2815	Binucleate	Unknown	Vicia faba	Valencia, Spain

TABLE 3. Anastomosis group, origin, size of ITS1, ITS2 and 5.8 s RDNA gene and EMBL accession number of tester isolates of *Rhizoctonia* spp.

Isolate	Anastomosis group	Isolated from	ITS 1	ITS2	Total	Accession No.
---------	-------------------	---------------	-------	------	-------	---------------

C-662	AG A	Soil	256	238	649	AJ242890
BN4	AG A (CAG2)	Soil	256	238	649	AJ242891
STR-2	AG B(o)	<i>Ipomoea batatas</i>	256	250	661	AJ242892
C-455	AG Bb	<i>Oryza sativa</i>	256	250	661	AJ242893
70B	AG C	Soil	257	243	655	AJ242894
C-610	AGD	unknown	25 8	243	656	AJ242895
Rhl55	AG E	unknown	25 8	243	656	AJ242896
C-653	AG G	unknown	252	252	659	AJ242897
AV-2	AG I	<i>Artemesia</i> sp.	251	255	661	AJ242898
184	AG J	unknown	251	255	661	AJ242899
FA59209	AG K	unknown	261	285	701	AJ242900
76146	AG L	unknown	259	278	692	AJ242901
TCI	AGN	unknown	254	285	694	AJ242902
76150	AG P	unknown	254	289	698	AJ242903
TC2	AG R	unknown	185	274	614	AJ242904
TC3	AG S	unknown	185	274	614	AJ242905

These isolates were kindly provided by M. Mazzola, USDA Agricultural Research Service, Tree Fruit Research Laboratory, 1104 N. Western Avenue, Wenatchee, WA 98801, U.S.A.

Typical *Rhizoctonia* isolates of this invention have been deposited under the Budapest Treaty, as follows:

isolate	deposit number	deposit date
<i>Rhizoctonia</i> (binucleate) Eab-F5	<i>Rhizoctonia</i> sp. CECT 20320	30 June 1999
<i>Rhizoctonia</i> (binucleate) Eab-F6	<i>Rhizoctonia</i> sp. CECT 20321	30 June 1999
<i>Rhizoctonia</i> (binucleate) Eab-S6	<i>Rhizoctonia</i> sp. CECT 20322	30 June 1999
<i>Rhizoctonia</i> (binucleate) Eab-S1	<i>Rhizoctonia</i> sp. CECT 20323	30 June 1999
<i>Rhizoctonia</i> (binucleate) Eab-F2	<i>Rhizoctonia</i> sp. CECT 20324	30 June 1999



The genomic DNA, asymmetric PCR and DNA sequencing procedures used in this study have been previously described. For DNA isolation, mycelia were crushed in liquid nitrogen, incubated at 65°C in extraction buffer and treated with RNase. Asymmetric PCR amplification, used to synthesize ssDNA, was done with a 50:1 molar ratio between the two primers (11). The primers used for amplification of the ITS regions were ITS4 and ITS5 (White *et al.*) and ITS1F and ITS4B (Gardes *et al.*). An automated thermal cycler (Perkin Elmer Cetus Corp. model 480, Norwalk, Connecticut) was used for the amplification reaction. The cycling parameters were initial denaturation (2.5 min at 94 °C) and 40 cycles of the following: denaturation, 15 s at 94 °C; annealing, 30 s at 55 °C when ITS1F/ITS4B were used as amplification primers (or 53 °C for ITS4/ITS5); and extension, 1.5 min at 72 °C.

In sequencing of the ITS-5.8 S rDNA region, PCR experiments carried out on the rDNA-ITS region of 30 isolates of BNR reproducibly amplified a single band. The amplification was terminated by a final extension for 10 min at 72 °C. PCR products were analyzed by electrophoresis in 1% agarose gels in TBE buffer, at constant voltage for two hours. Each PCR product was used for sequence determination after removing excess primers and nucleotides. Manual sequencing was performed using the Sanger's dideoxy method. All samples were sequenced at least twice in both directions. CLUSTALW alignment was used with Gap Open Penalty of 10.00 and Gap Extension Penalty of 0.05. The total length of the alignment obtained was 900 bp with 61 gaps. The length of the longest inferred gaps was 28 and 26 bp. The transition/transversion average was 1.764, calculated with DNAML from the PHYLIP package.

The nucleotide sequences determined are shown in Figure 1. They have been deposited in EMBL, and the given accession numbers are shown in the Tables.

The 5.8 S rDNA region was conserved among all the *Rhizoctonia* isolates examined and had a length of 155 bp. Among the isolates analysed, the sequence of ITS 2 was more divergent than the sequence of the ITS 1 region. Sequence homology among *Rhizoctonia* spp. isolates from Albacete varied from 90-99%, between isolates from Albacete and tester strains of binucleate anastomosis groups varied from 87-90 % and between Albacete isolates and multinucleate *R. solani* isolates varied from 68-88%.

For phylogenetic analysis, the sequence data were checked between complementary strands, and sequences were assembled using GCG Fragment Assembly System (Program Manual for the Wisconsin Package, version 8, Wisconsin). For the phylogenetic analysis we added sequences of multinucleate isolates from the different AGs of *Rhizoctonia solani* previously reported. These isolates were A-10 (AG 1), B60 (AG 2-2 IIIB), BC-10 (AG 2-2 IV), 1600 (AG 3)9 Pf-10 (AG 4 HG-1), Rh-165 (AG 4 HG-11), K31 (AG 5) HAMI-1 (AG 6), 1556 (AG 7), A68 (AG 8), V12M (AG 9) and TE2-4 (AG BI). Alignments of the ITS regions were performed using the multiple alignment program CLUSTALW (Thompson *et al.*) (IntelliGenetics, Inc., Mountain View, California), and are shown in the sequence listings. Only the sequences of the total ITS regions (including 5.8 S), without any nucleotide of the major ribosomal genes, were used for phylogenetic analysis.

Phylogenetic analysis of the aligned sequences was performed using the maximum parsimony analysis with the heuristic-search algorithm of the Phylogenetic Analysis Using Parsimony (PAUP) program 3.1.1. (Swofford *et al.*) with gaps treated as missing data. The heuristic search provides as optimal criteria maximum parsimony. The trees were rooted with the rDNA-ITS sequence of *Trichoderma harzianum* as outgroup. Data were resampled with 1000 bootstrap replicates by using the heuristic search of PAUP. The percentage of bootstrap replicates which yielded each grouping as a measure of statistical confidence was determined. Trees were confirmed from the distance matrix of Jukes & Cantor using the neighbor-joining method as it is in the program "Neighbor" from the PHYLIP package version 3.5. The Jukes & Cantor algorithm assumes that there are independent changes in all sites, with equal probability. The most parsimonious tree obtained by PAUP had a minimum length of 1212 steps, with consistency (CI) and retention (RI) indices of 0.606 and 0.751 respectively, and is shown in the Figure. In the tree, horizontal lengths in the branches indicate the relative genetic distances between the isolates, corresponding to the number of nucleotide substitutions.

Phylogenetic analysis separated all the *Rhizoctonia* spp. isolates from the *T. harzianum* outgroup isolate and clustered *Rhizoctonia* spp. isolates into two different groups, corresponding to binucleate and multinucleate isolates. All Spanish isolates from Albacete

except isolate S3 clustered differentiated from the binucleate tester isolates. The supposed AG 4 hypovirulent isolates 521 and Rh2815 clustered with the binucleate *Rhizoctonia* isolates. in the same group as the binucleate testers.

In this study, the first work up to date dealing with phylogeny of binucleate *Rhizoctonia*, the sequences of the ITS regions of 12 BNR isolates obtained in Spain, 16 BNR tester isolates, 2 previously identified as AG 4 hypovirulent isolates and 3 multinucleate isolates obtained in Spain have been determined and compared with known sequences of multinucleate *R. solani* strains obtained by Kuninaga *et al.*. The first conclusion clearly drawn upon observation of the resulting phylogenetic tree is that there are two separated branches, one grouping the multinucleate isolates and the other one including all the binucleate isolates, showing that multinucleate and binucleate isolates are genetically further from each other than multinucleate or binucleate isolates among themselves. This is supported by the fact that there is a cluster containing all the AGs of the multinucleate *Rhizoctonia*, which tend to be considered genetically isolated, whereas when a sufficiently high number of isolates is included in a study of this type, not only anastomosis groups, but also subgroups and ecological types are clearly separated.

The branch with binucleate isolates contains all the tester strains and the 12 BNR isolates from Spain. Surprisingly, in this branch were also included the supposedly multinucleate AG 4 hypovirulent strains 521 and Rh2815. Interestingly, in Boysen *et al.* these two isolates clustered together and apart from the seven AG 4 isolates studied and the AG 1 isolate used as an outgroup in the phylogenetic study. At that moment, the explanation given was that these two isolates clustered separately from the others based on pathogenicity, since all the other seven AG 4 and the one AG 1 isolates were highly virulent towards 5 different host plants (radish, cotton, onion, bean and carrot) and isolates 521 and Rh2815 were hypovirulent. It has also been shown that by using rDNA-ITS sequencing there is a clear grouping of the isolates according to pathogenicity and host range. Since the consistency of the tree was very high and the two branches clearly correlated with the number of nuclei of the isolates, it pointed to the possibility that strains 521 and Rh2815 had been misidentified. In fact, morphological criteria, such as cultural appearance, cannot be used to differentiate multinucleate from binucleate *Rhizoctonia*, since hyphal diameter is too similar and the formation of sclerotia or

melanine is also not discriminative. Therefore nuclear counting on isolates 521 and Rh2815 was performed, and showed that these isolates were BNR.

In the phylogenetic tree shown in this work, the molecular variation among *R. solani* isolates is bigger than among BNR isolates: it can be observed that the horizontal length of the branches of the tree, that represent genetic distance, are quite different for the multinucleate and the binucleate cluster. In the multinucleate cluster the branches are much longer than for the binucleate cluster, and the numbers above the internodes, which represent the bootstrap percentage from 1000 bootstrap replicates, are in general lower. This indicates that the ITS region of the binucleate isolates is much more similar among AGs than the same region of multinucleate isolates, and knowing that the ITS region is a good indicator of genetic evolution and relatedness, it can be concluded that binucleate isolates are more closely related among themselves than the multinucleate isolates.

For future work, from the sequences of Figure 1, it is possible to pick out sequences suited for use as PCR primers, as follows:

**Albacete**

**Eab Group**

CCCTATTAAGGGGCA

**Binucleated**

**Bin**

CCGTAAAAAAGTCTT

**Albacete**

**Ab+**

CCACACAAACCCATTGTATT

**Cornigerum**

**Corg**

AATTGTTCCCTTTGGGCAC

15

Albacete

Ab-2

GGAACCTTTATTGGAC

Albacete

Ab-3

CTCAATTTATTTTAAAACGA

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

TTGAATGAATATAGAGTTG

Eab-S2 and Eab-S4

TCGAATGAACAATGGGGATCG

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

TCCTCCGGGAG

Eab-S2 and Eab-S4

TTCTTCATTGAGG

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

AGACGGAGGACCGTAAAAAA

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

TTCCGTCTATTAAACC

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

ACACAAACCCCATTTGTATTTAAATTG

Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S5 and Eab-S6

ACATAAACTCCAATTTAATAAATCTG

Example 1: Initial Work

Pathogenicity tests were performed with the strains isolated from the field in Albacete on the following plant species: radish (*Raphanus sativus*), cauliflower (*Brassica oleracea* var. *botrytis*), onion (*Allium cepa*), turf grasses: zoysia grass (*Zoysia tenuifolia*), St. Augustine grass (*Stenotaphrum secundatum*) and Bermuda grass (*Cynodon dactylon*), tomato (*Lycopersicum esculentum*), carrot (*Daucus carota*) and cucumber (*Cucumis sativum*). Seeds were surface-disinfected in 1% sodium hypochlorite for 10 min. rinsed with sterile distilled water and aseptically blotted between two layers of wet sterile paper into petri dishes. They were allowed to germinate for 4 days at 26°C in the dark. Isolates were grown on water agar for 2 days at 26°C. Five germinated seeds were then placed around the fungal colony. The plates were incubated at room temperature in the lab bench, and pathogenicity was assayed for the first time 7-10 days after the tests were set, and subsequent observations were made afterwards every two days. The *Rhizoctonia solani* AG 4 strain Me8-2 (Boysen *et al.*, 1996) was used as positive control, since this strain is known to be very pathogenic to all plant species tested. A negative control of germinated seeds without fungus was included in each of the tests.

Crucifers (radish and cauliflower) are susceptible to all *Rhizoctonia* isolates tested, regardless of whether they were multinucleated or binucleate. The same was true for onion. Grass, tomato and carrot were susceptible to all multinucleated *Rhizoctoniae* (Me 8.2, aA1, aA2), but were not susceptible to any of the binucleate strains, therefore protection tests were done with these plant species. The strains randomly chosen to test protection from the pathogenic strain Me 8.2 were the binucleate strains aB and F2. Cucumber was susceptible to all strains tested, except for S7, therefore this strain was subsequently tested for protection from the pathogenic strain Me 8.2. The strain Me 8.2 showed significantly stronger pathogenicity than any other strains tested, and this was seen in shorter times needed to attack the plant, and complete killing of the seedling, whereas with other pathogenic strains the seedlings were damaged, but not as much affected as with isolate Me 8.2. In the pathogenicity tests was also observed that in some cases the seeds had not been completely surface sterilised, and in the negative control (no fungus) was observed growth of some fungi other than *Rhizoctonia*, yeast or bacteria, but when *Rhizoctonia* had colonised the seedlings, there was no growth of these contaminants.

The strains that did not show pathogenicity towards one or more plant species were tested for protection from the pathogenic strain Me 8.2. This was done by inoculating both the pathogenic strain and the supposedly protective strain. This was done in two ways, simultaneously (both strains inoculated at the same time) and successively (first the supposedly protective strain was allowed to colonise the plants surface for three days and then the pathogenic strain was inoculated). A negative control (no fungus), a positive control (the pathogenic strain Me 8.2), and the supposedly protective strain alone were included for each of the plant species in the tests. Protection was assayed for the first time 7 days after the tests were set, and subsequent observations were made every two days.

For grass, tomato and carrot, it was observed that the negative control (no fungus) presented contamination from other fungus, yeast and/or bacteria, whereas the control with protective strains B and F2 were clean, that is, *Rhizoctonia* did not allow any of the contaminants carried in the seed to grow. As expected, plants inoculated with protective strains B and F2 looked completely healthy, whereas plants inoculated with Me 8.2 were heavily injured. In the simultaneous test (that is, protective and pathogenic strains inoculated at the same time) there was no protection of the binucleate strains from the pathogenic Me 8.2. But in the successive test, in which the binucleate strains B and F2 were allowed to colonise the plant surface prior to adding the pathogenic strain, a very clear protective effect was observed, and the plants looked as healthy as the negative control and the control with the binucleate strains alone. Furthermore, it was also observed that the seedlings covered by the protective strains looked slightly bigger than the negative control, thus pointing to the possibility that there is an increased growth response induced by the non-pathogenic *Rhizoctonia* isolates. This has already been described by (Sneh *et al.*, 1986). In cucumber, although some protection was seen, we could not observe a clear protective effect of strain S7, therefore we could not conclude that strain S7 acts as a protective strain in cucumber.

Example 2: Detailed Investigations

Pathogenicity tests were performed to test whether the isolates candidates for protection against other fungal pathogens such as *Rhizoctonia solani* were themselves not pathogenic towards the plant species to be protected.

Seeds of each plant species to be tested were surface-disinfected in 1% sodium hypochlorite for 10 min, rinsed with sterile distilled water and aseptically blotted between two layers of wet sterile paper into petri dishes. They were allowed to germinate for 4 days at 26 °C in the dark. Isolates were grown in 9 cm diameter Petri dishes containing 2% water agar for 2 days at 26 °C. Five germinated seeds were then placed around the fungal colony. The plates were incubated at room temperature in the lab bench, and pathogenicity was assayed for the first time 7-10 days after the tests were set, and subsequent observations were made afterwards every two days. The *Rhizoctonia solani* AG 4 strain Me 8.2 (Boysen *et al.*) was always included as a positive control, since this strain is known to be very pathogenic to all plant species tested. A negative control of germinated seeds without fungus was included in each of the tests.

The strains that did not show pathogenicity towards one or more plant species were tested for protection from the pathogenic strain Me 8.2. This was done by inoculating both the pathogenic strain and the supposedly protective strain. This was done in two ways, simultaneously (both strains inoculated at the same time) and consecutively (first the supposedly protective strain was allowed to colonize the plants surface for three days and then the pathogenic strain was inoculated). A negative control (no fungus), a positive control (the pathogenic strain Me 8.2), and the supposedly protective strain alone were included for each of the plant species in the tests. Protection was assayed for the first time 7 days after the tests were set, and subsequent observations were made every two days. In some experiments, Eab-alphaB was also included. It is another experimental isolate employed for the purposes of comparison.

The following tests were carried out:



Protection test 1      Protection of grass, tomato and carrot seedlings with the binucleate *Rhizoctonia* isolates Eab-F2 and Eab-alphaB from the pathogenic *Rhizoctonia* Me 8.2 strain in simultaneous and consecutive tests in plates.

Protection test 2      Protection of pine using Eab-alphaB

Protection test 3      Protection of pine seedlings with the binucleate strains Eab-F2 and Eab-aB against the pathogenic strains Me 8.2 and Pin in simultaneous and consecutive tests under plate, flasks and greenhouse conditions.

Protection test 4      Protection of wheat seedlings and plants with binucleate *Rhizoctonia* strains against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests in plates, greenhouse and field conditions

Protection test 5      Protection of rape seedlings and plants with binucleate *Rhizoctonia* strains against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb, *Rhizoctonia* alphaB and *Altermaria* sp. in simultaneous and consecutive tests in plates and greenhouse conditions.

Protection test 6      Protection of alfalfa seedlings with the binucleate strains Eab-S1 and Eab-S7 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests under plate conditions.

Protection test 7      Protection of garlic bulbs with binucleate *Rhizoctonia* strains from pathogenic *Penicillium* sp. in direct inoculation assay at laboratory scale.

Protection test 8      Protection of sugar beet seedlings with the binucleate *Rhizoctonia* isolates Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-S1 and Eab-S7 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests in plates.

Protection test 9 Protection of onion seedlings with the binucleate *Rhizoctonia* isolates Eab-F1, Eab-F3, Eab-F5 and Eab-F6 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RA1b and *Fusarium* sp. in simultaneous and consecutive tests in plates.

All the tests have been performed in Petri dishes in the laboratory, except for the pine experiments, that have been done with pine seedlings growing on peat in a greenhouse in Albacete (Protection tests 2 and 3), and the wheat experiments, that have been performed in greenhouse as well as in field experiments (Protection test 4).

Data and graphs are included for providing a graphical view of the protection effects in the experiments. Data are referred in all the cases to the best protective strains concerning its behaviour in terms of protection. Thus, tables show the combination of those strains in each assay and the effects produced, expressed as coefficients using Butler's scale (1 to 5, being 5 the more pathogenic effect) for disease measurement. Finally, those data are expressed in bar histograms.

#### Protection test 1

Protection of grass, tomato and carrot seedlings with the binucleate *Rhizoctonia* isolates Eab-F2 and Eab-alphaB from the pathogenic *Rhizoctonia* Me 8.2 strain in simultaneous and consecutive tests in plates.

PLANT SPECIES	grass, tomato, carrot
STRAINS TESTED	Eab-F2, Eab-alphaB
TIMING	simultaneous, consecutive
DATE	November 1997

Two protective isolates, Eab-F2 and Eab-alphaB, were used to assay protection effects in grass, tomato and carrot against the pathogenic *Rhizoctonia* isolate Me 8.2. The pathogenic strain and the supposedly protective strain were inoculated in two ways, simultaneously (both strains inoculated at the same time) and consecutively (first the supposedly protective strain

was allowed to colonize the plants surface for three days and then the pathogenic strain was inoculated). A negative control (no fungus), a positive control (the pathogenic strain Me 8.2), and the supposedly protective strain alone were included for each of the plant species in the tests.

The tables shows the results of plate assays. Presence or absence of protection effects is indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts: 1. 17/11/97, 2. 19/11/97. 3. 21/11/97. Date of simultaneous inoculation: 11/11/97: date of consecutive inoculation: 14/11/97

#### Grass, tomato and carrot protection test

	C- (no fungus)	C+ Me 8.2	Eab-F2	Eab- alphaB	Eab-F2/ Me 8.2 Simult.	Eab-F2/ Me 8.2 consec.	Eab- alphaB/ Me 8.2 simult.	Eab- alphaB/ Me 8.2 consec.
Tomato	0-0-0	4-4-5	0-0-0	0-0-0	4-4-5	0-0-0	4-4-5	0-0-0
Carrot	0-0-0	5-5-5	0-0-0	1-1-1	5-5-5	1-1-1	4-5-5	0-1-1
Grass	0-0-0	0-2-3	0-0-0	0-0-0	0-1-2	0-0-0	0-3-3	0-0-0

#### Disease scale (BUTLER CODE) (referred to infected surface)

0. without detectable infection, 1. infection until 10 %., 2. infection 11-30 %. 3. infection 31-50 %, 4. infection 51-80 %, 5 more than 80 %

In all three plant species, grass, tomato and carrot, there was a very clear protection effect of the two binucleate strains tested, Eab-F2 and Eab-alphaB, from the pathogenic isolate Me 8.2 when the protective strain was inoculated and allowed to colonize the surface before the pathogenic strain was applied, that is. in the consecutive essay. In the simultaneous essay, the pathogenic strain Me 8.2 showed the same pathogenicity behaviour when applied together with the protective strain that when applied alone. That is probably because the pathogenic strain grows faster than the protective strain, therefore is the first one to colonize the plant surface.

It was also observed that in the positive control (no fungus) there were contaminations produced by other fungi, yeast and bacteria. whereas the plates inoculated only with the protective strain showed no growth of such contaminants.

#### Protection test 2

PLANT SPECIES Pine  
STRAINS TESTED Eab-alphaB  
DATE April 1998

The initial material were trays containing each one 54 pots. In each pot there was a variable number of seeds, since more than one seed had been sowed per pot. At the moment of inoculation of the strains, seedlings had not emerged yet. The pathogenic strain used was Pin, a strain initially isolated from pine. The protective strain was Eab-alphaB.

We tried changing different parameters. such as inoculum concentration of the protective strain, the protective strain was added as a suspension in liquid medium or grown on sterile sand and adding different fungicide concentrations.

The seedlings had not emerged at the moment when we applied the protective strain. The observations after applying the strains were hindered by the fact that the substrate used to grow the pine seedlings had a very high sand content, and the humidity compacted the sand, creating a hard layer that did not allow emergence of the seedlings. Therefore it was not possible to draw any conclusions from this experiment, and it was repeated with seedlings that were already emerged (see Protection Test 3)

#### Protection test 3

Protection of pine seedlings with the binucleate strains Eab-F2 and Eab-aB against the pathogenic strains Me 8.2 and Pin in simultaneous and consecutive tests under plate, flasks and greenhouse conditions.

PLANT SPECIES Pine  
STRAINS TESTED Eab-F2, Eab-aB  
DATE May 1998

PLATES AND FLASKS. We have tested 2 binucleate *Rhizoctonia* protective strains (Eab-F2 and Eab-aB) against two pathogenic *Rhizoctonia* strains, Me 8.2 and Pin. The strain Me 8.2 has been chosen as a pathogenic control, due to the fact that this pathogenic strain has been shown to have a broad range of hosts (including pine). Pin strain has been isolated as pathogen on pine in the field. Protective strains were inoculated in culture flasks and petri dishes (assays were duplicate under these two systems) with solid media (water agar: 15 g Bacto Agar, 11 distilled water), containing one (in the case of flasks) or two (petri dishes) pine seedling each one. Pine seeds were previously sterilized and germinated according to the following protocol: introduce seeds 36 h on distilled water at 5 °C; then, wash the seeds 5 min. with a solution containing 3 drops of Tween 80 on 100 ml of distilled water; wash again with distilled water (at least three washes); introduce seeds 15 min on a solution containing hydrogen peroxide 30%; wash (three times at least) with distilled water; put 15-20 seeds on a petri dish containing water agar (15 g Bacto Agar, 11 distilled water) and incubate them inverted at 21 °C on the dark about 14 days. The pathogenic strains were applied at two moments with respect to the protective strains, simultaneously or consecutively. 7 days after simultaneous inoculation in order to determine whether some kind of advantage in terms of accession and colonization of the host plant could lead to an increasing protection effect.

GREENHOUSE. The initial plant material were pine seedlings that were already emerged in greenhouse trays containing each one 54 pots. In each spot there was a variable number of seedlings, since more than one seed had been sowed per pot. At the moment of inoculation of the fungus, the seedlings were about 2 cm above the soil surface. We tested two supposedly protective strains, Eab-F2 and Eab-aB, two supposedly pathogenic strains, Me 8.2 and Pin. and two ways of applying the protective strain, directly in liquid suspension in the soil, or the strain

previously grown in sterile peat. We studied protection at two different moments: simultaneous, by applying the protective and the pathogenic strain at the same moment, and consecutive, by allowing the protective strain to colonize the plant surface and applying the pathogenic strain three weeks after the protective strain. As an additional test, we also treated some trays with fungicide. The first observations were done 6 weeks after the application of the pathogenic strains, and subsequent observations were done also 9, 12 and 24 weeks after.

There were a total of 18 trays, and each of the trays was treated in a different way:

1. negative control, no treatment
2. protective strain Eab-aB in liquid suspension
3. protective strain Eab-F2 in liquid suspension
4. protective strain Eab-aB in peat
5. protective strain Eab-F2 in peat
6. pathogenic strain Me 8.2
7. pathogenic strain Pin
8. negative control, the same media of the fungal suspension
9. simultaneous test with the protective Eab-aB and the pathogenic Me 8.2
10. simultaneous test with the protective Eab-aB and the pathogenic Pin
11. simultaneous test with the protective Eab-F2 and the pathogenic Me 8.2
12. simultaneous test with the protective Eab-F2 and the pathogenic Pin
13. consecutive test with the protective Eab-aB and the pathogenic Me 8.2
14. consecutive test with the protective Eab-aB and the pathogenic Pin
15. consecutive test with the protective Eab-F2 and the pathogenic Me 8.2
16. consecutive test with the protective Eab-F2 and the pathogenic Pin
- 2.1. fungicide
- 2.2. fungicide

PLATES. Due to the nature and design of plate and flask protection tests in pine seedlings, results are expressed as visual estimation of photographs.

GREENHOUSE. The countings of the surviving pine seedlings, performed 24 weeks after inoculation of the pathogenic strains, were as follows:

1.	negative control. no treatment	122
2.	protective Eab-alphaB in liquid suspension	150
3.	protective Eab-F2 in liquid suspension	120
8.	negative control, the same media of the fungal suspension	94
7.	pathogenic strain Pin	9
10.	simultaneous with protective Eab-alphaB and pathogenic Pin	5
12.	simultaneous with protective Eab-F2 and pathogenic Pin	43
14.	consecutive with protective Eab-alphaB and pathogenic Pin	119
16.	consecutive with protective Eab-F2 and pathogenic Pin	104
2.1.	fungicide	102
2.2.	fungicide	92

Applying statistics, the results can be interpreted as follows:

PLATES. Results from both flasks and plates tests have shown a strong correspondence between laboratory and greenhouse results. In this sense, a clear protection effect can be observed by the two protective strains assayed (Eab-F2 and Eab-aB) against pin and Me 8.2 strains. As observed at greenhouse scale (see comments below), this protection effect is increased when consecutive inoculation of the pathogenic strain is made.

GREENHOUSE. We have tested two protective strains, Eab-F2 and Eab-aB, and two pathogenic strains, Me 8.2 and Pin. The pathogenic strains were added at the same time than the protective strain (simultaneous test) and 3 weeks after the protective strain was added (consecutive test).

It was observed that:

- the pathogenic strain Pin was strongly pathogenic (see "Pin" at the photographs), whereas the strain Me 8.2, which is a very severe pathogen in other plant species, was not pathogenic towards pines.
- the protective strains inoculated alone were not pathogenic at all towards the pine seedlings (see "Eab-aB and Eab-F2")
- there was no difference in applying the protective strain in liquid suspension or in peat

- there was no protection effect of Eab-aB against Pin when they were applied at the same time (simultaneous test), but there was some protection effect of F2 against Pin at the simultaneous test.
- there was a very clearly observed protection effect of Eab-aB and Eab-F2 against Pin if they were inoculated some weeks before the pathogenic strain (consecutive test), therefore allowed to colonize the substrate and carry out their protection effect. This protection effect was very high, and the number of living seedling was of the same range as for the negative controls (no fungus) or the protective strain alone.

#### Protection test 4

Protection of wheat seedlings and plants with binucleate *Rhizoctonia* strains against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests in plates, greenhouse and field conditions

PLANT SPECIES wheat

#### STRAINS TESTED

PROTECTIVES (PLATES): 2815, 521, Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-F6, Eab-F7, Eab-S1, Eab-S2, ab-S3, Eab-S4, Eab-S5, Eab-S6, Eab-S7. (GREENHOUSE): Eab-F5, Eab-S6.

(FIELD): Eab-F5, Eab-S6.

PATHOGENS (PLATES): *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb, *Rhizoctonia* aB, *Fusarium*, *Acremonium*.

(GREENHOUSE): *Fusarium*, *Rhizoctonia* RAlb.

(FIELD): *Fusarium*, *Rhizoctonia* RAlb.

TIMING PLATES: simultaneous, consecutive

GREENHOUSE: simultaneous, consecutive seedling stage, consecutive plant stage

FIELD: simultaneous, consecutive seedling stage, consecutive plant stage.

DATE January-April 1999



## 4a. PLATES

We have tested 15 binucleate *Rhizoctonia* protective strains (described in the introduction) against five pathogenic fungi; three of them were *Rhizoctonia* strains, including *Rhizoctonia* Me 8.s as a positive control, one *Rhizoctonia* RAlb strain isolated from wheat debris in Albacete and a third *Rhizoctonia*, aB, isolated from alfalfa in Albacete. The other two pathogens assayed (*Fusarium* and *Acremonium* strains) were also isolated from wheat samplings in Albacete. Assays were carried out in petri dishes containing about 20 ml of water-agar (15 g Bacto Agar, 1l distilled water) and performed at room temperature. Seeds were previously surface sterilized (ethanol 70% 2'; sodium hypochlorite 5% with Tween 0.005% 15' and at least five washes in distilled water) and germinated in culture chamber at 28°C in the dark.

Pathogenic strains were applied at two moments with respect to the protective fungi, in order to determine whether some kind of advantage in terms of colonization of the host by the protective strain could lead to an increasing protection effect. Thus, in some cases binucleate *Rhizoctonias* were inoculated simultaneously with the pathogen, and in others, pathogenic strains were inoculated 3 days after the protective ones.

The table shows the results of plate assays. Presence or absence of protection effects are indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts: 1. 29/01/99, 2. 01/02/99, 3. 03/02/99. Date of simultaneous inoculation: 20/01/99; date of consecutive inoculation: 23/01/99

## Wheat protection test

	Me 8.2 Simult	Me 8.2 Conse c.	RAlb Simult	RAlb. Conse c.	Alpha B Simult	Alpha B Conse cutive	Fus. Simult	Fus. Conse c.	Acrem Simult	Acrem Conse c.
None	3-3-3		1-2-3		0-0-0		1-3-3		0-0-0	
2815	0-0-0	0-0-0	0-0-1	1-1-1	0-0-0	0-0-0	0-0-0	0-0-0	1-1-1	0-1-1
521	0-0-0	0-0-0	1-1-2	1-2-3	0-0-1	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0

Eab-F1	1-2-2	0-0-0	1-2-3	1-1-2	0-0-0	0-1-1	0-2-2	0-0-0	0-0-0	0-0-0
Eab-F2	0-0-1	0-2-2	2-2-3	0-0-0	0-0-0	0-0-0	1-3-3	0-0-0	0-0-0	1-1-1
Eab-F3	1-1-2	0-0-0	1-1-1	0-1-1	0-0-0	0-1-1	1-2-3	0-0-0	0-0-0	0-0-0
Eab-F5	0-0-0	0-0-1	0-1-2	0-0-1	0-0-0	0-0-0	1-2-3	0-0-0	0-0-0	0-0-0
Eab-F6	1-1-1	0-0-0	1-2-3	0-1-2	0-0-0	0-0-0	1-2-3	0-0-0	0-0-0	0-0-0
Eab-F7	1-1-1	0-0-0	1-2-3	1-1-2	1-1-1	0-0-0	1-1-1	0-0-0	0-0-0	0-1-1
Eab-S1	0-0-0	1-2-2	1-2-2	1-2-3	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0
Eab-S2	0-0-0	0-1-1	1-2-3	1-1-2	0-0-0	0-0-0	1-1-2	0-0-0	0-0-0	0-0-0
Eab-S3	1-2-3	0-1-1	1-2-2	2-3-3	0-0-0	0-0-0	1-1-2	0-0-0	0-0-0	0-0-0
Eab-S4	0-0-0	0-0-0	0-1-3	1-1-1	0-0-0	0-0-0	1-2-2	0-0-0	0-0-0	0-0-0
Eab-S5	0-0-0	0-1-1	0-1-3	0-1-2	0-0-0	0-0-0	1-2-3	0-0-0	0-0-0	0-0-0
Eab-S6	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0
Eab-S7	1-1-2	0-1-2	1-2-3	1-2-3	0-1-1	0-0-0	1-2-3	0-0-0	0-0-0	0-0-0
A1	2-3-3	1-3-3	2-3-3	2-2-3	1-1-1	2-3-3	2-3-4	2-3-3	2-3-3	2-3-3
A2	2-3-3	2-2-3	1-2-3	2-3-4	1-1-1	2-3-3	2-3-3	2-3-3	2-3-3	2-3-3

In general terms, protection effects are increased when consecutive inoculation of first the protective strain and then the pathogenic strain is done. Thus, when pathogenic strain is

inoculated time after, protective fungi have the advantage in colonizing the host plant, increasing significantly the protection.

Plate assays shows that the degree and effectiveness of the protection depends on the type of pathogenic strain assayed, as well as the protective strain tested. In this way, it has been shown that protection of wheat from pathogens is clearly more effective if *Fusarium* and *Rhizoctonia* Ralb are inoculated as pathogens. Plants inoculated with pathogenic *Rhizoctonia* Me 8.2 strain, (used as a positive control since this isolate is known to be highly pathogenic) were less protected and this little protection effect was only seen with consecutive inoculation.

Finally, strains Eab-S6 and Eab-F5 have been proven to be the most interesting protective strains at laboratory assay. Those two strains showed the highest rates of protection against the pathogens tested and were selected to perform greenhouse and field assays. In addition, we decided to select *Fusarium* and *Rhizoctonia* RAlb as the most suitable pathogenic fungi to be tested in greenhouse and field together with the protective fungi above mentioned because they are isolated from the same location where greenhouse and field tests were going to be carried out.

#### 4b. GREENHOUSE

The two protective strains selected, S6 and F5 have been shown to have highest protection properties against two selected pathogenic strains, *Fusarium* and *Rhizoctonia* RAlb. Greenhouse assay was carried out in 48-pot trays disposed over a cement layer and grouped in two separated blocks (sterilized vs non-sterilized peat), distant 2 m from each others, with a separation between trays inside each block of at least 50 cm in order to prevent crossed contaminations. Three wheat seeds were disposed in each pot to reduce the existence of non-germinating ones; after early emergence of the seedlings, some of them were removed and one seedling per pot was left. The trays (32 study cases) were inoculated at three moments: simultaneous inoculation of the protective and pathogenic strains 20 days after sowing; consecutive, adding the pathogen at seedling stage (14 days after protective strain inoculation) and consecutive at mature plant stage, adding the pathogen 30 days after protective strain inoculation. Factors such as presence or absence of fungicide (fungicide treatment was a

combination of Metil-Toclofos and Tiram. effective against *Fusarium*-like and *Rhizoctonia* diseases), as well as the use of sterilized or unsterilized peat were introduced in order to evaluate the effect of such variables on the protection. An scheme showing the several statistical cases and interactions considered is provided below.

1	control	simultaneous	control	no	no
2	control	plant	<i>Rhizoctonia</i>	simultaneous	no
3	control	simultaneous	<i>Fusarium</i>	no	simultaneous
4	control	plant	<i>Rhizoctonia</i>	no	simultaneous
5	<i>Rhizoctonia</i>	plantlet	control	simultaneous	no
6	<i>Rhizoctonia</i>	simultaneous	<i>Fusarium</i>	no	no
7	<i>Rhizoctonia</i>	plantlet	control	no	simultaneous
8	<i>Rhizoctonia</i>	plantlet	<i>Rhizoctonia</i>	simultaneous	simultaneous
9	<i>Rhizoctonia</i>	simultaneous	control	no	simultaneous
10	<i>Rhizoctonia</i>	simultaneous	<i>Fusarium</i>	simultaneous	simultaneous
11	control	plant	control	no	no
12	control	plant	control	simultaneous	simultaneous
13	<i>Rhizoctonia</i>	plantlet	<i>Rhizoctonia</i>	no	no
14	control	plantlet	control	no	no
15	<i>Rhizoctonia</i>	plant	control	simultaneous	no
16	control	simultaneous	control	no	no
17	<i>Rhizoctonia</i>	simultaneous	control	simultaneous	no
18	<i>Rhizoctonia</i>	simultaneous	control	simultaneous	no
19	<i>Rhizoctonia</i>	plant	<i>Fusarium</i>	no	no
20	control	simultaneous	<i>Rhizoctonia</i>	simultaneous	no
21	control	simultaneous	control	simultaneous	simultaneous
22	<i>Rhizoctonia</i>	plant	control	no	simultaneous
23	<i>Rhizoctonia</i>	simultaneous	<i>Rhizoctonia</i>	simultaneous	simultaneous
24	<i>Rhizoctonia</i>	simultaneous	control	no	simultaneous
25	control	plantlet	<i>Fusarium</i>	simultaneous	no
26	control	plantlet	control	simultaneous	simultaneous

27	<i>Rhizoctonia</i>	simultaneous	<i>Rhizoctonia</i>	no	no
28	control	simultaneous	<i>Rhizoctonia</i>	no	simultaneous
29	control	simultaneous	control	simultaneous	simultaneous
30	control	plantlet	<i>Fusarium</i>	no	simultaneous
31	<i>Rhizoctonia</i>	plant	<i>Fusarium</i>	simultaneous	simultaneous
32	control	simultaneous	<i>Fusarium</i>	simultaneous	no

Greenhouse tests will be considered as finished when wheat plants will be completely mature and ready to harvest. At the moment, measurement of crop efficiency is not yet finished and results are based on comparative visual estimations (those are shown as several figures at the end).

<i>Rhizoctonia</i> S6	simultaneous	<i>Fusarium</i>	control
control	simultaneous	control	<i>Fusarium</i>
<i>Rhizoctonia</i> F5	simultaneous	<i>Rhizoctonia</i> AG2	<i>Fusarium</i>
control	plant	control	control
control	plantlet	<i>Fusarium</i>	control
<i>Rhizoctonia</i> S6	simultaneous	control	<i>Fusarium</i>
<i>Rhizoctonia</i> F5	simultaneous	control	control
<i>Rhizoctonia</i> F5	plantlet	control	control
<i>Rhizoctonia</i> S6	plant	<i>Rhizoctonia</i> AG2	control
control	simultaneous	<i>Rhizoctonia</i> AG2	control
control	plant	control	<i>Fusarium</i>
control	simultaneous	control	control
<i>Rhizoctonia</i> F5	plant	<i>Fusarium</i>	<i>Fusarium</i>
control	simultaneous	<i>Fusarium</i>	<i>Fusarium</i>
<i>Rhizoctonia</i> S6	plantlet	control	<i>Fusarium</i>
control	plantlet	<i>Rhizoctonia</i> AG2	<i>Fusarium</i>

Protection against the two pathogenic strains used, *Fusarium* sp and *Rhizoctonia* Raib (Albacete strain) were observed (visual estimation). This protection effect was specially patent against *Fusarium* sp., where both protective strains Eab-S6 and Eab-F5 were shown to

be remarkably efficient in protecting and preventing *Fusarium* disease. Symptoms of *Fusarium* attack appeared by middle of May (five months after inoculation) and consisted on a progressive drying (the so-called "añublo de espigas") of the spike just before pollination, becoming sterile many of them. Finally, correlation between visual estimates and crop efficiency (determined by statistical methods) is expected; This would provide an effective method to quantify protection.

#### 4c. FIELD

The same protective and pathogenic strains used in greenhouse were performed in field conditions. Strains were inoculated in 10 x 1.7 m plots at the same three moments as in greenhouse. Presence or absence of fungicide (same combined treatment used) were introduced as additional variables. A scheme showing the statistical cases and interactions considered is provided below.

At the moment, we have only available results from laboratory and greenhouse designs, since field test are currently developing.

#### Protection test 5

Protection of rape seedlings and plants with binucleate *Rhizoctonia* strains against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RA1b, *Rhizoctonia* alphaB and *Alternaria* sp. in simultaneous and consecutive tests in plates and greenhouse conditions.

PLANT SPECIES rape

STRAINS TESTED

PROTECTIVES (PLATES): 2815, 521, Eab-F1, Eab-F3, Eab-F6, Eab-S1, Eab-S4, Eab-S6, Eab-S7,

(GREENHOUSE): Eab-F6, Eab-S6.

PATHOGENS (PLATES): *Rhizoctonia* Me 8.2 strain, *Rhizoctonia* RA1b strain, *Rhizoctonia* ???strain, *Alternaria*

(GREENHOUSE):*Alternaria*. RAlb.

#### TIMING

PLATES: simultaneous, consecutive

GREENHOUSE: simultaneous, consecutive seedling stage, consecutive plant stage

DATE January-April 1999

#### 5a. PLATES

9 protective strains were used to assay protection effects in rape seedlings against pathogenic fungi. In addition, 3 pathogenic fungi were used in the protection assay; two of them were *Rhizoctonia* strains, including *Rhizoctonia* Me 8.2 (as a pathogenic control) and one *Rhizoctonia* RAlb isolated from Albacete. The last pathogen assayed (an *Alternaria* sp. strain) was also isolated from rape samplings in Albacete. Assays were carried out in petri dishes containing about 20 ml of water-agar (15 g Bacto Agar, 1l distilled water) and performed at room temperature. Seeds were previously surface sterilized (ethanol 70% 2'; sodium hypochlorite 5% with Tween 0.005 % 15' and at least five washes in distilled water) and germinated in culture chamber at 28°C in the dark.

Pathogenic strains were applied at two moments with respect to the protective fungi in order to determinate whether some kind of advantage in terms of colonization of the host by the protective strain could lead to an increasing protection effect. Thus, in some cases binucleate *Rhizoctonias* were inoculated simultaneously with the pathogen, and in others, pathogenic strains were inoculated 2 days after the protective ones.

Table shows the results of plate assays. Presence or absence of protective effects is indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts realized on: 1. 29/01/99, 2. 01/02/99, 3. 03/02/99. Date of simultaneous inoculation: 20/01/99; date of consecutive inoculation: 22/01/99

Rape protection test

	Me 8.2 Consec.	Me 8.2 Simult.	AlphaB Consec.	AlphaB Simult.	Altern. Consec.	Altern. Simult.	RAIb Consec.	RAIb Simult.
None	4-5-5		2-3-5		1-1-3		3-5-5	
2815	0-0-0	2-3-3	0-1-1	0-0-0	0-2-2	0-0-0	0-0-0	2-2-3
521	1-2-2	3-4-5	0-0-0	0-0-0	0-0-0	0-0-1	0-1-1	1-4-5
Eab-F1	0-0-0	5-5-5	0-0-1	1-2-3	0-0-1	0-0-0	0-0-0	1-3-3
Eab-F3	1-1-2	3-5-5	1-1-2	1-2-3	0-1-1	0-0-0	0-0-0	3-4-5
Eab-F6	0-0-0	3-4-4	0-0-0	1-2-3	0-0-0	0-0-0	0-0-0	1-3-3
Eab-S1	0-0-0	4-5-5	0-0-0	0-1-2	0-0-0	0-0-0	0-0-0	3-4-5
Eab-S4	0-1-1	3-5-5	0-1-1	1-2-2	0-0-1	0-0-1	0-0-0	1-3-3
Eab-S6	0-0-1	0-1-2	1-1-2	1-2-2	0-1-2	0-0-1	0-0-0	0-0-0
Eab-S7	0-0-1	3-5-5	0-0-0	2-3-3	0-0-0	0-0-1	0-0-0	1-4-4
A1	2-4-5	3-4-5	2-4-4	1-4-4	2-4-5	1-3-4	1-4-4	2-4-4
A2	3-4-4	1-4-5	1-4-4	2-3-4	1-4-5	2-3-4	2-4-5	2-3-4

After finishing the plate assays, the best two protective strains were selected in order to test them at greenhouse scale. We have observed a remarkably protection effect by binucleate *Rhizoctonia* strains against the selected pathogens in rape.

*Rhizoctonia* Me 8.2 strain, *Rhizoctonia* RAIb and *Alternaria* sp. were demonstrated to be pathogens to rape and protection against them was observed in laboratory. This was specially patent for those plates with consecutive inoculation moment. where protection effects were well observed since the firsts stages of the assay.

Finally, the two protective strains chosen were Eab-S6 and Eab-F6 (they result always protective against all pathogens tested in consecutive inoculation and, in some cases, even at simultaneous moment), and were assayed in greenhouse against *Rhizoctonia* RAIb and *Alternaria* strain as selected pathogens (these last were selected because of the origin of the strain, being isolated from rape plants in the study area).

#### 5b. GREENHOUSE



The two protective strains selected, Eab-S6 and Eab-F6 have been shown to have highest protection against two selected pathogens (*Alternaria* and *Rhizoctonia* RA1b). Greenhouse assay was carried out in 48-pot trays disposed over a cement layer and grouped in two separated blocks (sterilized vs non-sterilized peat), distant 2 m from each others, with a separation between trays inside each block of at least 50 cm in order to prevent crossed contaminations. The trays (32 study cases) were inoculated at three moments: simultaneous inoculation of the protective strains and pathogens 7 days after sowing; consecutive, adding the pathogen at seedling stage (9 days after simultaneous inoculation) and consecutive at mature plant stage, adding the pathogen 21 days after simultaneous inoculation). Factors such as presence or absence of fungicide, as well as the use of sterilized or unsterilized peat as substrate were introduced in order to evaluate the effect of such variables on the protection effects. Temperature inside the greenhouse and irrigation system were also controlled and periodically recorded. A scheme showing the several statistical cases and interactions considered for further statistic studies is provided below.

1	control	simultaneous	control	no	no
2	control	plant	<i>Rhizoctonia</i>	simultaneous	no
3	control	simultaneous	<i>Fusarium</i>	no	simultaneous
4	control	plant	<i>Rhizoctonia</i>	no	simultaneous
5	<i>Rhizoctonia</i>	plantlet	control	simultaneous	no
6	<i>Rhizoctonia</i>	simultaneous	<i>Fusarium</i>	no	no
7	<i>Rhizoctonia</i>	plantlet	control	no	simultaneous
8	<i>Rhizoctonia</i>	plantlet	<i>Rhizoctonia</i>	simultaneous	simultaneous
9	<i>Rhizoctonia</i>	simultaneous	control	no	simultaneous
10	<i>Rhizoctonia</i>	simultaneous	<i>Fusarium</i>	simultaneous	simultaneous
11	control	plant	control	no	no
12	control	plant	control	simultaneous	simultaneous
13	<i>Rhizoctonia</i>	plantlet	<i>Rhizoctonia</i>	no	no
14	control	plantlet	control	no	no
15	<i>Rhizoctonia</i>	plant	control	simultaneous	no
16	control	simultaneous	control	no	no
17	<i>Rhizoctonia</i>	simultaneous	control	simultaneous	no

18	<i>Rhizoctonia</i>	simultaneous	control	simultaneous	no
19	<i>Rhizoctonia</i>	plant	<i>Fusarium</i>	no	no
20	control	simultaneous	<i>Rhizoctonia</i>	simultaneous	no
21	control	simultaneous	control	simultaneous	simultaneous
22	<i>Rhizoctonia</i>	plant	control	no	simultaneous
23	<i>Rhizoctonia</i>	simultaneous	<i>Rhizoctonia</i>	simultaneous	simultaneous
24	<i>Rhizoctonia</i>	simultaneous	control	no	simultaneous
25	control	plantlet	<i>Fusarium</i>	simultaneous	no
26	control	plantlet	control	simultaneous	simultaneous
27	<i>Rhizoctonia</i>	simultaneous	<i>Rhizoctonia</i>	no	no
28	control	simultaneous	<i>Rhizoctonia</i>	no	simultaneous
29	control	simultaneous	control	simultaneous	simultaneous
30	control	plantlet	<i>Fusarium</i>	no	simultaneous
31	<i>Rhizoctonia</i>	plant	<i>Fusarium</i>	simultaneous	simultaneous
32	control	simultaneous	<i>Fusarium</i>	simultaneous	no

Greenhouse assays are actually developing and results have been referred just to laboratory tests. No field assays were disposed for rape because of the weather conditions by this period of the year.

#### Protection test 6

Protection of alfalfa seedlings with the binucleate strains Eab-S1 and Eab-S7 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RA1b and *Fusarium* sp. in simultaneous and consecutive tests under plate conditions.

PLANT SPECIES alfalfa

STRAINS TESTED Eab-S1, Eab-S7

TIMING simultaneous, consecutive

DATE May 1999

Two protective binucleate *Rhizoctonia* strains (Eab-S1 and Eab-S7) were used to assay protection effect in alfalfa seedlings against selected pathogenic fungi. In this sense, 3 pathogenic fungal strains were assayed in the experimental design: two of them were *Rhizoctonia* strains, including *Rhizoctonia* Me8.2 (as a pathogenic control) and *Rhizoctonia* RAlb strain isolated from Albacete. The third pathogen assayed (*Fusarium* strain) was isolated from alfalfa debris in the same province. Assays were carried out at laboratory scale in petri dishes containing about 20 ml of water-agar (15 g Bacto Agar, 1l distilled water) and performed at room temperature. Seeds were previously surface sterilized (ethanol 70% 2'; sodium hypochlorite 5% with Tween 0.005% 15' and at least five washes in distilled water) and pre-germinated in culture chamber at 28°C in the dark. The pathogenic strains were applied at two moments with respect to the protective strains, simultaneously or 6 days after in order to determine whether some kind of advantage in terms of accession and colonization of the host plant could lead to an increasing protection effect.

Tables shows the results of plate assays. Presence or absence of protection effects is indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts: 1. 06/05/99, 2. 10/05/99, 3. 13/05/99. Date of simultaneous inoculation: 27/04/99; date of consecutive inoculation: 03/05/99

#### Alfalfa protection test

1*	Me 8.2 simult.	Me 8.2 consec.	RAlb simult.	RAlb consec.	Fus. simult.	Fus. consec.
None	1-2-4		2-3-5		1-1-3	
Eab-S1	0-1-2	0-0-0	1-3-4	0-0-0	1-2-2	0-1-1
Eab-S7	2-3-4	1-1-1	2-4-4	0-0-0	1-1-2	0-0-0

2*	Me 8.2 simult.	Me 8.2 consec.	RAlb simult.	RAlb consec.	Fus. simult.	Fus. consec.
None	1-3-5		2-5-5		1-2-3	
Eab-S1	1-2-2	0-0-0	2-3-4	0-0-0	0-0-1	0-0-0
Eab-S7	3-3-4	0-1-1	2-4-5	0-0-0	1-1-1	0-0-2

\*1 and 2 means duplicate experiments

Plant and protective strains controls

PLANT (-)	0	0	0
Eab-S1	0	0	0
Eab-S7	1	2	2

After finishing the plate assays, it can be observed that protection effect is produced by the two protective strains (Eab-S1 and Eab-S7) used against the three pathogens assayed. In general terms, protection is increased when consecutive inoculation of first the protective strains and then the pathogenic strain is made. Thus, when pathogenic strain is inoculated time after, protective fungi seems to have the advantage in terms of competition for colonization sites in the tissues of the host plant, increasing significantly the protection. This phenomenon is especially patent when both *Rhizoctonia* Me 8.2 and *Rhizoctonia* RAlb are present as pathogens. In such cases, protection effect is always increased when consecutive inoculation of the protective strain is made.

#### Protection test 7

Protection of garlic bulbs with binucleate *Rhizoctonia* strains from pathogenic *Penicillium* sp. in direct inoculation assay at laboratory scale.

PLANT SPECIES    garlic

STRAINS TESTED   Eab-F1, Eab-F7, Eab-S4, Eab-S6

TIMING           simultaneous direct inoculation

DATE               May 1999

We have tested 4 binucleate *Rhizoctonia* protective strains (above mentioned) against one pathogenic *Penicillium* sp. strain. We have chosen this fungal strain due to the fact that this pathogenic fungus has been usually reported as a garlic pathogen and furthermore, it was always present in all the garlic bulbs used for this study, causing conspicuous damages, specially in roots. Due to the nature of the pathogen, we decided to apply protective strains with a direct inoculation method. Thus, protective strains were firstly incubated in liquid culture (CM medium: malt extract 5 g, yeast extract 5 g, Glucose 5 g., 1l distilled water) and after 5 days, fungal biomass was recovered and then, inoculated in culture flasks with solid media (water agar: 15 g Bacto Agar, 1l distilled water) directly over non-sterilized garlic bulbs naturally infected with *Penicillium* sp.

Table shows the results of flasks assay. Presence or absence of protection effect is indicated as estimation of infected vegetal surface (using Butler scale). Two numbers correspond to 2 counts: 1. 10/05/99, 2. 14/05/99. Next to the scale number, a sign + between brackets means increment of biomass. Date of inoculation: 07/05/99.

#### Garlic protection test

	Eab-S4	Eab-S6	Eab-F1	Eab-F7
1	0 (++)	0 (+)	0 (++)	0
2	0 (++)	0 (+)	0 (++)	0

After finishing the flasks assays, it can be observed some protection effect against the pathogen. In this sense, direct inoculation of protective strain seems to prevent and limit *Penicillium* effects. Though infection by *Penicillium* is usually located and restricted at the base of the bulb colonizing the young roots, we have observed that infection spreads under wet and warm conditions (as in the case of flask cultures) invading the seedlings and even blocking sometimes germination of the bulbs. In general terms, all the protective *Rhizoctonias* tested, prevented the spread of *Penicillium* sp. and in some cases (Eab-F1 and Eab-S7), increasing of vegetal biomass was clearly observed with the presence of protective strains.

## Protection test 8

Protection of sugar beet seedlings with the binucleate *Rhizoctonia* isolates Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-S1 and Eab-S7 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests in plates.

PLANT SPECIES    sugar beet  
STRAINS TESTED   Eab-F1, Eab-F2, Eab-F3, Eab-F5, Eab-S1 and Eab-S7  
TIMING            simultaneous, consecutive  
DATE              May-June 1999

6 protective isolates were used to assay protection effects in sugar beet seedlings against selected pathogenic fungi. In this sense, 3 pathogenic fungal strains were used in experimental design; two of them were *Rhizoctonia* strains, including *Rhizoctonia* Me 8.2 (as a pathogenic control) and one *Rhizoctonia* RAlb strain isolated from Albacete. The third pathogen assayed (*Fusarium* sp. strain) was isolated from sugar beet debris in the same locality. Assays were carried out at laboratory scale in petri dishes containing about 20 ml of water-agar (15 g Bacto Agar, 1l distilled water) and performed at room temperature. Seeds were previously surface sterilized (ethanol 70% 2'; sodium hypochlorite 5% with Tween 0.005% 15' and at least five washes in distilled water) and pre-germinated in culture chamber at 28 °C in the dark. The pathogenic strains were applied at two moments with respect to the protective strains, simultaneously or 8 days after simultaneous inoculation in order to evaluate the effects on the protection of these two moments.

Tables shows the results of plate assays. Presence or absence of protection effects are indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts: 1. 01/06/99, 2. 03/06/99, 3. 06/06/99. Date of simultaneous inoculation: 20/05/99; date of consecutive inoculation: 28/05/99

Sugar beet protection test

	Me 8.2 simult.	Me 8.2 consec.	RAlb simult.	RAlb consec.	Fus. simult.	Fus. consec.
None	4-4-5		3-3-4		1-2-3	
Eab-F1	4-5-5	0-1-1	5-5-5	0-0-0	1-3-3	0-0-0
Eab-F2	5-5-5	2-4-4	5-5-5	0-1-1	3-4-4	1-1-1
Eab-F3	2-3-5	4-4-4	5-5-5	0-0-0	3-4-4	2-3-3
Eab-F5	4-5-5	3-4-4	5-5-5	4-5-5	4-5-5	4-4-4
Eab-S1	2-3-4	3-4-4	4-5-5	0-0-0	2-2-3	0-0-0
Eab-S7	3-4-5	0-0-0	5-5-5	0-0-0	2-3-3	1-1-1

In general terms, protection effects are increased when consecutive inoculation of first the protective strain and then the pathogenic strain is done. Thus, when pathogenic strain is inoculated time after, protective fungi have the advantage in colonizing the host plant tissues, increasing significantly the protection.

Plate assays shows that the degree and effectiveness of the protection effect depends on the type of pathogenic strain assayed, as well as the protective strain tested. In this way, it has been shown that protection of sugar beet from pathogens is clearly more effective if *Rhizoctonia* Ralb is inoculated as pathogen (excepting for the Eab-F5 strain). Plants inoculated with pathogenic *Rhizoctonia*. Me 8.2 strain, were in general terms less protected (only strains Eab-F1 and Eab-S7 showed a clear protection effect) and this little effect was always seen with consecutive inoculation of the pathogen. On the other hand, plants inoculated with *Fusarium* sp. strain were protected by 4 of the 6 strains (Eab-F1, Eab-F2, Eab-S2 and Eab-S7) but, as in the case of Me 8.2 strain, only in the plates where consecutive inoculation was done.

#### Protection test 9

Protection of onion seedlings with the binucleate *Rhizoctonia* isolates Eab-F1, Eab-F3, Eab-F5 and Eab-F6 against the pathogenic strains *Rhizoctonia* Me 8.2, *Rhizoctonia* RAlb and *Fusarium* sp. in simultaneous and consecutive tests in plates.

PLANT SPECIES onion  
 STRAINS TESTED Eab-F1, Eab-F3, Eab-F5, Eab-F6  
 TIMING simultaneous, consecutive  
 DATE June 1999

4 protective isolates were used to assay protection effects in onion seedlings against selected pathogenic fungi. In this sense, 3 pathogenic fungal strains were used in experimental design: two of them were *Rhizoctonia* strains, including *Rhizoctonia* Me 8.2 (as a pathogenic control) and one *Rhizoctonia* RAlb strain isolated from Albacete. The third pathogen assayed (*Fusarium* sp. strain) was isolated from onion debris in the same locality. Assays were carried out at laboratory scale in petri dishes containing about 20 ml of water-agar (15 g Bacto Agar, 1 l distilled water) and performed at room temperature. Onion seeds were previously sterilized and pre-germinated according the following protocol: introduce seeds on a solution of H<sub>2</sub>O<sub>2</sub> 30% 1h; then wash (3-4 times at least) with sterile water; dry seeds and put them on petri dishes (40-50 seeds per dish) and incubate inverted at 27 °C in the dark for two days. The pathogenic strains were applied at two moments with respect to the protective strains, simultaneously or 5 days after simultaneous inoculation in order to evaluate the effects on the protection of these two moments.

Tables shows the results of plate assays. Presence or absence of protection effects are indicated as estimation of infected vegetal surface (using Butler scale). Three numbers correspond to 3 counts: 1. 15/06/99, 2. 17/06/99, 3. 19/06/99. Date of simultaneous inoculation: 09/06/99; date of consecutive inoculation: 14/06/99

specially for Eab-F6 strain Onion protection test

	Me 8.2 simult.	Me 8.2 consec.	RAlb simult.	RAlb consec.	Fus. simult.	Fus. consec.
None	3-4-5		3-4-5		2-4-5	
Eab-F1	0-0-0	0-0-1	0-1-1	0-0-0	0-0-0	0-0-1
Eab-F3	0-0-1	0-0-0	0-1-1	0-0-1	0-0-0	0-0-0



43

Eab-F5	0-1-2	0-0-0	0-0-1	0-0-0	0-0-0	0-0-0
Eab-F6	0-0-1	0-0-0	0-0-0	0-0-0	0-0-0	0-0-0

## Plant and protective strains controls

PLANT (-)	0	0	0
Eab-F1	0	1	1
Eab-F3	0	0	1
Eab-F5	0	0	0
Eab-F6	0	0	0

In general terms, protection effects are observed for all the strains assayed in both consecutive and simultaneous inoculation moments, although protection is more effective when consecutive inoculation is made. Regarding the protective strains, all the binucleate *Rhizoctonias* showed a clear protection effect, specially for Eab-F6 strain.

Laboratory scale plate assays resume.

S t r a i n s	WHEAT PATHOGENS						ALFALFA PATHOGENS					
	Me 8.2		Fus. p5		RALb		Me 8.2		Ralb		Fus. p5	
	S	C	S	C	S	C	S	C	S	C	S	C

44

Eab	++	+	++	++	++	++	+	++	-	++	+	++
-S1												
Eab	++	++	+	++	++	++						
-S6												
Eab	+	+	-	++	-	++						
-F2												
Eab	++	+	-	++	+	++						
-F5												
Eab	+	++	+	++	-	+						
-F6												
Eab	++	++	-	++	-	+						
-S4												
Eab	+	+	-	++	-	-	-	+	-	++	+	++
-S7												
Eab	-	++	-	++	-	-						
-F1												
Eab												
-aB												
Eab	+	++	+	++	-	-						
-F7												

(-): No protection

S: Simultaneous inoculation

(+: Protection

C: Consecutive inoculation

(++): Good protection

ID: Direct inoculation

S	RAPE PATHOGENS	GA	CARROT PATHOGENS
		RLI	
		C	

45

t r a i n s	Me 8.2		Altern. p13		RAIb		<i>Peni cilli um</i>	Me 8.2		RAIb	
	S	C	S	C	S	C	I.D.	S	C	S	C
Eab -S1	-	++	+	++	-	+					
Eab -S6	+	+	++	+	+	++	++	+	+	-	++
Eab -F2								-	+		
Eab -F5											
Eab -F6	-	++	+	+	+	++					
Eab -S4	-	+	+	+	-	++	++				
Eab -S7	-	+	+	++	-	++					
Eab -F1	-	++	++	++	-	++	++				
Eab -aB								-	+	+	+
Eab -F7							+				

(-): No protection

S: Simultaneous inoculation

(+: Protection

C: Consecutive inoculation

(++): Good protection

ID: Direct inoculation

S t r a i n s	TOMATO PATHOG ENS		GRASS PATHOG ENS		ONION PATHOGENS						PINE PATHOGENS			
	Me 8.2		ME 8.2		Me 8.2		Fus. p10		RALb		pin		Me 8.2	
	S	C	S	C	S	C	S	C	S	C	S	C	S	C
Eab -S1														
Eab -S6														
Eab -F2	-	++	+	++							-	++	+	++
Eab -F5					+	++	++	++	+	++				
Eab -F6					+	++	++	++	++	++				
Eab -S4														
Eab -S7														
Eab -F1					++	+	++	+	+	++				
Eab -aB	-	++	-	++							+	++	+	++

47

Eab					+	++	++	++	+	+				
-F3														
Eab														
-F7														

(-): No protection

S: Simultaneous inoculation

(+: Protection

C: Consecutive inoculation

(++): Good protection

ID: Direct inoculation

## REFERENCES

Boysen. M., Borja, M., del Moral, C., Salazar, O. and Rubio. V. (1996) Identification at strain level of *Rhizoctonia solani* AG4 isolates by direct sequence of asymmetric PCR products of the ITS regions Curr. Genet., 29. 174-181.

Cubeta. M. A., Echandi, E., Abernethy, T. and Vilgalys. R. (1991) Characterization of anastomosis groups of binucleate *Rhizoctonia* species using restriction analysis of an amplified ribosomal RNA gene Phytopathology, 81. 1395-1400.

Duncan. S., Barton. J. E. and O'Brien. P. A. (1993) Analysis of variation in isolates of *Rhizoctonia solani* by random amplified polymorphic DNA assay Mycol. Res., 97, 1075-1082.

Felsenstein, J. (1993) PHYLIP manual version 3.5. Washington. University of Washington.

Gardes. M., and Bruns, T. D. (1993) ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts Molec. Ecol. 2:113-118

Harris, A. R., Schisler. D. A., Neate, S., and Ryder, M. H. 1993. Suppression of damping off caused by *Rhizoctonia solani* and growth promotion in bedding plants by binucleate *Rhizoctonia* spp. Soil. Biol. Biochem. 25:263-268.

- Herr, L. J. 1995. Biological control of *Rhizoctonia solani* by binucleate *Rhizoctonia* spp. and hypovirulent *R. solani* agents. Crop Protection 14:179-186.
- Jabaji-Hare, S. H., Meller, Y., Gill, S. and Charest, P. M. (1990) Investigation of genetic relatedness among anastomosis groups of *Rhizoctonia solani* using cloned DNA probes Can. J. Plant Pathol., 12, 393-404.
- Julián, M. C., Dullemans, A. M., Silfhout, C., and Keijer, J. 1997. Nuclear behavior in homokaryotic and heterokaryotic fruiting of *Thanatephorus cucumeris* (*Rhizoctonia solani*) anastomosis group 1, subgroup IC. Mycologia 89:361-374.
- Kataria, H. R. and Gisi, U. (1989) Recovery from soil and sensitivity to fungicide of *Rhizoctonia solani* Mycol. Res., 92, 458-462.
- Keijer, J., Houterman, P. M., Dullemans, A. M. and Korsman, M. G. (1996) Heterogeneity in electrophoretic karyotype within and between anastomosis groups of *Rhizoctonia solani* Mycol. Res., 100, 789-797.
- Kimura, M. (1980) A simple method for estimating evolutionary rates through comparative studies of nucleotide sequences. J. Mol. Evol. 16, 111-120.
- Kuninaga, S. and Yokosawa, R. (1980) A comparison of DNA base compositions among anastomosis groups in *Rhizoctonia solani* Kühn Ann. Phytopath. Soc. Japan, 46, 150-158.
- Kuninaga, S. and Yokosawa, R. (1982) DNA base sequence homology in *Rhizoctonia solani* Kühn II. Genetic relatedness within anastomosis group 2 Ann. Phytopath. Soc. Japan, 48, 668-673.
- Kuninaga, S. and Yokosawa, R. (1982) DNA base sequence homology in *Rhizoctonia solani* Kühn. I. Genetic relatedness within anastomosis group 1 Annu. Phytopath. Soc. Japan, 48, 659-667.

Kuninaga, S. and Yokosawa, R. (1984) DNA base sequence homology in *Rhizoctonia solani* Kühn. IV. Genetic relatedness within AG-4 Annals. Phytopathol. Soc. Japan. 50. 322-330.

Kuninaga, S. and Yokosawa, R. (1985) DNA base sequence homology in *Rhizoctonia solani* Kühn. V. Genetic relatedness within AG-6 Annu. Phytopathol. Soc. Japan. 50. 346-352.

Kuninaga, S. and Yokosawa, R. (1985) DNA base sequence homology in *Rhizoctonia solani* Kühn. VI. Genetic relatedness among seven anastomosis groups Annals Phytopathological Society of Japan, 51, 127-132.

Kuninaga, S. and Yokosawa, R. (1985) DNA base sequence homology in *Rhizoctonia solani* Kühn. VII. Genetic relatedness between AG-BI and other anastomosis groups Annu. Phytopathol. Soc. Japan. 51. 133-138.

Poromarto, S. H., Nelson. B. D., and Freeman. T. P. 1998. Association of binucleate *Rhizoctonia* with soybean and mechanism of biocontrol of *Rhizoctonia solani*. Phytopathology 88:1056-1067

Saitou, N. and M.. Nei. (1987) The neighbour-joining method: a new method for reconstructing phylogenetic trees Molecular Biology and Evolution. 4. 406-425.

Sanger, F., Nicklen. S. and Coulson, A. R. (1977) DNA sequencing with chain-termination inhibitors Proc. Natl. Acad. Sci. USA, 74. 5463-5467.

Sneh, B., Zeidan, M., Ichielevich-Auster, M., Barash, I. and Koltin, Y. (1986) Increased growth responses induced by a nonpathogenic *Rhizoctonia solani* Can. J. Bot.. 64. 2372-2378.

Sneh, B., Jabaji-Hare, S., Neate. S., and Dijst, G.. eds. 1996. *Rhizoctonia* Species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Sneh, B. 1998. Use of non-pathogenic or hypovirulent strains to protect plants against closely related fungal pathogens. *Biotechnol. Adv.* 16:1-23.

Sneh, B., and Ichielevich-Auster, M. 1998. Induced resistance of cucumber seedlings caused by some non-pathogenic *Rhizoctonia* (np-R) isolates. *Phytoparasitica* 26:27-38.

Swofford, D. L. (1991) PAUP: Phylogenetic Analysis Using Parsimony, version 3.1.1. Champaign, IL., Illinois Natural History Survey.

Thompson, J. D., Higgins, D. G. and Gibson, T. J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice *Nucl. Acids Res.* 22, 4673-4680.

Vilgalys, R. and Gonzalez, D. (1990) Ribosomal DNA restriction fragment length polymorphisms in *Rhizoctonia solani* *Phytopathology*, 80, 151-158.

White, T. J., Bruns, T. D., Lee, S., and Taylor, J. W. (1990) Amplification and direct sequencing of fungal ribosomal RNA for phylogenetics pages 315-322 in: *PCR protocols: A guide to Methods and Applications*. M. A. Innis, D. H. Gelfand, J. J. Sninsky, T. J. White, eds. Academic Press. San Diego.



## CLAIMS

1. A method for the protection of plants from pathogenic fungi which comprises biocontrol by the use of a binucleate *Rhizoctonia* strain selected by molecular detection of the ITS ribosomal sequence.
2. A method according to claim 1 where the plant is selected from tomato, pepper, cucumber, lettuce, radish, parsley and other herbs, bean, potato, sugar beet, carrot, garlic, onions, alfalfa, grass, wheat, rape, pine and other trees, as well as flowers.
3. A method according to claim 1, where the plant is a field crop, container-grown crop, water bed crop, vegetable or fruit crop, ornamental crop, nursery crop, garden crop, forest crop, or greenhouse crop.
4. A method according to claim 1 wherein the binucleate *Rhizoctonia* strain is selected using the ITS1 data.
5. A method according to claim 4, wherein the strain has an ITS1 sequence which is at least 90% homologous with one of SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, or SEQ ID NO:28.
6. A method according to claim 5, wherein the strain is selected from *Rhizoctonia* sp. CECT 20320, *Rhizoctonia* sp. CECT 20321, *Rhizoctonia* sp. CECT 20322, *Rhizoctonia* sp.

CECT 20323, *Rhizoctonia* sp. CECT 20324, *Rhizoctonia* sp. ATCC 64643. and *Rhizoctonia* sp. CECT 2815.

7. A binucleate *Rhizoctonia* strain which has an ITS1 sequence which includes a fragment of at least 10 nucleic acid bases which is present in any of one of SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, or SEQ ID NO:28.

8. A concentrate for use in plant protection, containing viable material of a binucleate *Rhizoctonia* strain according to claim 7.

9. A method for the protection of plants from pathogenic fungi which comprises biocontrol by the use of a binucleate *Rhizoctonia* strain according to claim 7.

10. A primer for use in identifying binucleate *Rhizoctonia* strains of use in biocontrol, the primer being selected from SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, and SEQ ID NO: 14.

1/64

Eab-F2 .....AGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-F3 .....TT.....AGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-F5 .....GG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-S5 .....T.....CG..A..TA..AA..GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-S6 .....GGG.A..TA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-F1 .....TT.....AGAGGGGAA.TA..AAAAGTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-F6 .....TTTAGAGGG.AAGTA..AAAAGTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-F7 ..TTCCCTT..T..TTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-S1 .....A.....GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-S2 .....GTCGTAACAAGGT.TTCCGT.AGGT  
 Eab-S4 .....GTCGTAACAAGGT.TTCCGT.AGGT

521 .....  
 Eab-S3 .....A..AA..GTCGTAACAAGGT.TTCCGT.AGGT  
 RH2815 .....  
 AGK ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGN ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGP ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGBO GCT.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGBB GCT.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGJ ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGG ..T.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGC ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGA ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 CAG2 ....C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGD ..T.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGE ....C..TTGGTCATTTAGAGG..AAGTATTA...GTCGTAACAAGGT.TTCCGT.AGGT  
 AGI ..T.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGL ..T.C..TTGGTCATTTAGAGG..AAGTA..AAA.GTCGTAACAAGGT.TTCCGT.AGGT  
 AGR .....  
 AGS .....

Eab-A1 .....A..AA..GTCG.AACAAGGT.TTCCGT.AGGT  
 Eab-A2 .....A..AA..GTCG.AACAAGGT.TTCCGT.AGGT  
 Eab-S7 .....A..A...GTCGTAACAAGGT.TTCCGT.AGGT  
 68AG8 .....  
 K31AG5 .....  
 16001AG3 .....  
 TPV12MAG9 .....  
 HG1HAM6 .....  
 TE25AGBI .....  
 1556AG7 .....  
 HG1PF10 .....  
 HGIIRH165 .....  
 AG22IV .....  
 AG22IIB .....  
 1AAG1 .....  
 Tharzianum .....T.....A.G.A..A...GTCGTAACAAGGTCT.CCGTT.GGT

Fig. 1(A)

SUBSTITUTE SHEET (RULE 26)

09744502-20544250

09/744502

2/64

Eab-F2 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-F3 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-F5 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-S5 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-S6 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-F1 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-F6 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-F7 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-S1 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT.A...T...AGAG.....T.T...G..GT.TG  
 Eab-S2 GAA.CCT.GCGGAAGG.ATCATT...T.CGAA.TG....AA.CAA..T....G.G.GG...AT...CG..GT.TG  
 Eab-S4 GAA.CCT.GCGGAAGG.ATCATT...T.CGAA.TG....AA.CAA..T....G.G.GG...AT...CG..GT.TG  
  
 521 .....A...G....A.....G.....T....CG..GT.TG  
 Eab-S3 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.C...TT...AGAG.....T....CG..GT.TG  
 RH2815 .....A.....G....AG.....T.TT..G..GT.TG  
 AGK GAA.CCT.GCGGAAGG.ATCATT...TT.GAA...CC...A..C...TT...G.GAG....T....CG..GT.TG  
 AGN GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGP GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.CA.C.TT...G.GAG....T....CG..GT.TG  
 AGBO GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.C...TT...AGAG.....T....CG..GT.TG  
 AGBB GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.C...TT...AGAG.....T....CG..GT.TG  
 AGJ GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGG GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AAT...GT...AGAG.....T.T...G..GT.TG  
 AGC GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGA GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 CAG2 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGD GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGE ..AAACCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...AGAG.....T.T...G..GT.TG  
 AGI GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA..A.C.T...G.GAG....T.TT..G..GT.TG  
 AGL GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.CA..T....G.GAG....T.TT..G..GT.TG  
 AGR ....C...GCGGA.GG..TC.TT.....AA.TG....AAT...G....A.A.AGG.....GA.GT.TG  
 AGS ....C...GCGGA.GG..TC.TT.....AA.TG....AAT...G....A.A.AGG.....GA.GT.TG  
  
 Eab-A1 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.T...T..AAT...GT...AGAG.....T.TT..G..GT.TG  
 Eab-A2 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.T...T..AAT...GT...AGAG.....T.TT..G..GT.TG  
 Eab-S7 GAA.CCT.GCGGAAGG.ATCATT...TT.GAA.TG....AA.C...TT...G.GAG....T....CG..GT.TG  
 68AG8 .....AA...T..TT.....AA.TG....AA....G....AG.....T.T...G..GT.TG  
 K31AG5 .....AA...T..TT.....AA.TG....A.....G....G.G.G...ATATTT.G..GT.TG  
 16001AG3 .....AA...T..TT...T...AA.TG....AA....G....AG.....TTT.G..GT.TG  
 TPV12MAG9 .....AA...T..TT...T...AA.TG....A.....G....GAG.....TTT.G..GT.TG  
 HG1HAM6 .....AA...T..TT.....AA.TG..T..A.....G....AG.....TT..G..GT.TG  
 TE25AGBI .....AA...T..TT...TT..AA.TG....AA....G....AG.....TTT.G..GT.TG  
 1556AG7 ..AA.....TTA.....AA.TG..T.....G....GAG.....TT..G..GT.TG  
 HG1PF10 GAA...T.....T...AA.TG..T..A.....G....AG.....TTT.G..GT.TG  
 HGIIRH165 GAA...T.....T...AA.TG..T..A.....G....AG.....TTT.G..GT.TG  
 AG22IV ..AA...T.....TTA.....AA...C...AA..A..G....G.G....TATTTG..GT.TG  
 AG22IIIB ..AA...T.....TT..AAA..C...AA..A..G....G.G....TATTTG..GT.TG  
 1AAG1 .....TTA..TT..AA.TG....A.....G....GAG.....T.T...GA.GT.TG  
 Tharzianum GAA.CC.AGCGGA.GGGATCATTACC...GA...G..TTTA..CAAC.T.CCCA.A.A.CCC.....AA.TGTG

Fig. 1(B)  
SUBSTITUTE SHEET (RULE 26)

09744502-071301

09/744502

3/64

Eab-F2 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-F3 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-F5 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-S5 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-S6 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-F1 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-F6 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-F7 TC...GCTGGC..T...CC....TC...C.....GG.GAG...C..A..TGTGCAC  
 Eab-S1 TC...GCTGGC..T...CC....TC..GC.....GG.GAG...C..A..TGTGCAC  
 Eab-S2 TC...GCTGGC..TT..C....T.T..CA.....TT.....G...AGG...C..A..TGTGCAC  
 Eab-S4 TC...GCTGGC..TT..C....T.T..CA.....TT.....G...AGG...C..A..TGTGCAC  
  
 521 TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 Eab-S3 TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 RH2815 T.A..GCTGG...T...CC....CT...AA..TTAA.C.TT.....GGGGGC..A..TGTGCAC  
 AGK TC...GCTGGC..T....G...TCTT..A...T.....TT.....GGC.AG...C..A..TGTGCAC  
 AGN TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 AGP TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 AGB0 TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 AGBB TC...GCTGGC..T....G...TCTT.....T.....GGC.AG...C..A..TGTGCAC  
 AGJ TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.C..A..TGTGCAC  
 AGG TC...GCTGGC.....CC....T.TT.CA.C.T.....GGGG..TA..TGTGCAC  
 AGC TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.C..A..TGTGCAC  
 AGA TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.C..A..TGTGCAC  
 CAG2 TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.C..A..TGTGCAC  
 AGD TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.C..A..TGTGCAC  
 AGE TC...GCTGGC.....CC....CTT.CA.C.T.....GGGG.CC.A..TGTGCAC  
 AGI T...TGCTGGC.....CC....CTT.....T.A...TT.....GGGG.C..A..TGTGCAC  
 AGL T...TGCTGGC.....CC....CTT.....T.A...TT.....GGGG.C..A..TGTGCAC  
 AGR ..A.TGCTGGC.....C..AGGTCT..C...T.....GG...C..A..TGTGCTC  
 AGS ..A.TGCTGGC.....C..AGGTCT..C...T.....GG...C..A..TGTGCTC  
  
 Eab-A1 T.A..GCTGGC..T...C....CT...AA..TTAAAC.TT.....GGGGGC..A..TGTGCAC  
 Eab-A2 T.A..GCTGGC..T...C....CT...AA..TTAAAC.TT.....GGGGGC..A..TGTGCAC  
 Eab-S7 TC...GCTGGC..T....G...TCTT..A...T.....TT.....GGC.AG...C..A..TGTGCAC  
 68AG8 T.A..GCTGG...T...CC.A..T.T...AA..TT.....T.....GGG..C..AA.TGTGCAC  
 K31AG5 T.A..GCTGGC..T...C..A..TC...AA..TT.....T.....GGG..C..A..TGTGCAC  
 16001AG3 T.A..GCTGGCC.....C..A..T.T...AA..TT.....T.....AGG...C..A..TGTGCAC  
 TPV12MAG9 T.A..GCTGGCC.T.....A..T.T..CA...TT.....T.....GGG..C..A..TGTGCAC  
 HGIHAM6 T.A..GCTGGCC.....C.GA..T.TT..AA..T.A...TTT.....GG...G...C..A..TGTGCAC  
 TE25AGBI T.A..GCTGG...T...CC.A..T.T...AA..T.....TT.....GG...G...C..AAC.GTGCAC  
 1556AG7 T.A..GCTGGCC.TT...GA..T.T..CA...T.....TTT..C.TA.....GG...C..A..TGTGCAC  
 HG1PF10 T.A..GCTGG...T...CC....CT...AA..TTAA.C.TT.....GG...GGG..C..A..TGTGCAC  
 HGIIRH165 T.A..GCTGGC..T...CC....T....AA..TTAAAC.TT.....GGGGGC..A..TGTGCAC  
 AG22IV T.A..GCTGGC..T...CC.A..T.T...A.....GTTT.....GG..AG...C..A..TGTGCAC  
 AG22IIIB T.A..GCTGGC..T...CC.A..T.T...A.....GTTT.....GG..AG...C..A..TGTGCAC  
 1AAG1 T...TGCTGGCC.TTT.....TCT...A.CCTTAA...TTTGGC..A.GG..AGGGGGC..A..TGTGCAC  
 Tharzianum ..AA.....CCAT..ACC.A.....AAC.T....GTT.G.CCT.CGGC..GGG.....GT.CAC

Fig. 1(C)

SUBSTITUTE SHEET (RULE 26)

FOET 20" 20544260

09/744502

4/64

Eab-F2 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-F3 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-F5 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-S5 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-S6 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC.G...TGTGCA.C.T...TGTG  
 Eab-F1 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-F6 .AC...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-F7 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-S1 G.C...T.TT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 Eab-S2 G.CC..T.TT...C.TCT.ATTT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 Eab-S4 G.CC..T.TT...C.TCT.ATTT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
  
 521 G.CC..T.T...C.TCT..TTT...CA..TCC...A...CACA.C.C..CC..TGTGCA.C.T...TGTG  
 Eab-S3 G.CC..T.T...C.TCT.ATT...CA..TCC...A...CACA.C.C..CC..TGTGCA.C.T...TGTG  
 RH2815 .ACC..T.TT...C.TCT..TT....CA..TCCC..AT.A.CACA.C.C.....TGTGCA.CCT....GTG  
 AGK G.CC..T.T...C.TCT..TTT...CA..TCC...A...CACA.CAC..CCC.TGTGCA.C.T...TGTG  
 AGN G.CC..T.T...C.TCT..TTT...CA..TCC...A...CACA.C.C..CC..TGTGCA.C.T...TGTG  
 AGP G.CC..T.T...C.TCT.ATT...CA..TCC...A...CACA.C.C..CC..TGTGCA.C.T...TGTG  
 AGBO G.CC..T.T...C.TCT.ATT...CA..TCC...A...CACA.C.C..CC..TGTGCA.CCT....GTG  
 AGBB G.CC..T.T...C.TCT.ATT...CA..TCC...A...CACA.C.C..CC..TGTGCA.CCT....GTG  
 AGJ G.CC..T.T...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 AGG G.CC..T.T...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 AGC G.CC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 AGA G.CC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T.CCC...G  
 CAG2 .CCC.T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T.CCC...G  
 AGD G.CC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 AGE G.CC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 AGI .ACC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 AGL .ACC..T.....CCTCT.ATT...CA..TCC...A...CACA.CAC..C..TGTG.AAC.T...TGTG  
 AGR G.C...T.T...CC...GA.....CAA.TC....TTATC.CAT.ACA.CC..TGTGCA.C..AC..TGTG  
 AGS G.C...T.T...CC...GA.....CAA.TC....TTATC.CAT.ACA.CC..TGTGCA.C..AC..TGTG  
  
 Eab-A1 .ACC..T.TT...C.TCT..TT....CA..TCCC..AT.A.CACA.C.C.....TGTGCA.CCT....GTG  
 Eab-A2 .ACC..T.TT...C.TCT..TT....CA..TCCC..AT.A.CACA.C.C.....TGTGCA.CCT....GTG  
 Eab-S7 G.CC..T.T...C.TCT..TTT...CA..TCC...A...CACA.CAC..CCC.TGTGCA.C.T...TGTG  
 68AG8 .ACC..T.....CCTCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.CC....TGTG  
 K31AG5 .ACC..TAT...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.C.T...TGTG  
 16001AG3 .ACC..T.TT...C.TCT..TT....CA..TCCC..AT.A.CACA.C.C.....TGTG.AAC.T...TGTG  
 TPV12MAG9 .ACC..T.TT...C.TCT..TT....CA..TCCC..A...CACA.CAC..C..TGTG.AACC....TGTG  
 HG1HAM6 .ACC..T.T...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.CC....TGTG  
 TE25AGBI .ACC..T.TT...C.TCT..TT....CA..TCC...AT.A.CACA.CAC..C..TGTG.AACC....TGTG  
 1556AG7 .ACC..T.T...C.TCT..TT....CA..TCC...A...CACA.CAC..C..TGTGCA.CC....TGTG  
 HG1PF10 .ACC..T.TT...C.TCT..TT....CA..TCCC..AT.A.CACA.C.C.....TGTGCA.CC....TGTG  
 HGIIRH165 .ACC..T.TT...C.TCT..TT....CA..TCCC..ATA..CACA.C.C.....TGTGCA.CC....TGTG  
 AG22IV .ACC..T.TTTG..C.TCT..TTTTTT.AA.TCC...A...CACA.C.C..CC..TGTG.AACC....TGTG  
 AG22IIIB .ACC..T.TTTG..C.TCT..TTTTTT.AA.TCC...A...CACA.CAC..C..TGTG.AACC....TGTG  
 1AAG1 .ACC..T.T...C.TCT..TT....CA..TCC...AT...CACA.C.C..CCC.TGTGCA.C.T...TGTG  
 Tharzianum G.CCCC.....GGG..T..G.....C..GTC...G.....CA.....GCCCC.G.G.AACC.A...G.G

Fig. 1(D)

SUBSTITUTE SHEET (RULE 26)

FOET 20 20544250

Eab-F2 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-F3 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-F5 AGAC..GG.A.G.G.ACC..G.T.....AA...AAA...TGTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-S5 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-S6 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-F1 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-F6 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-F7 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-S1 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.C.C.G..T.....C.....T...ATT..  
 Eab-S2 AGAC..GG.A..T..A...G.T.....A.GT.AAAAAA.GTC..TT.C...G..T.....C.....T.G..T..C..  
 Eab-S4 AGAC..GG.A.G.G.ACC..G.T.....AA...AAAA...GTC..TT.CT..G..T.....C.....T.....

521 AGACT.GG.A.G.G..CC..G.T.....AA...AAA...G.CC.TT.C..AG..T.....C.....T.GC.T..  
 Eab-S3 AGACT.GG.A.G.G..CC..G.T.....AA...AAA...G.CC.TT.C..AG..TG...C.....T...A..  
 RH2815 AGAC.AG..A..TG.....TTTT.CTA.G..A.....G.....GG.GG...AAG.G.....  
 AGK AGACT.GG.A.G...ACC.GGT.....AA...AAAAA...GTC..TT.C..AG..T.....C.....T.GC....  
 AGN AGACTAGG.A.G.G..CC..G.T.....AA...AAA...G.C..TTTC..AG..T.....C.....T.GC....  
 AGP AGACT.GG.A.G.G..CC..G.T.....AA...AAA...G.CC.TT.C..AG..T.....C.....T.GC....  
 AGB0 AGACT.GG.A.G...CCC.G.T.....AA...AAA...G.CC.TT.C..AG..T.....C.....T.GC....  
 AGBB AGACT.GG.A.G...CCCC..T.....AA...AAA...G.CC.TT.C..AG..T.....C.....T.GC....  
 AGJ AGAC..GG.A.G.GG.C....TTT...AA.TTA.....GTC..TT.C.C.G..T.....C.....T...AT..  
 AGG AGAC..GG.A.G.GG.C....TTT...AA.TTA.....GTC..TT.C.C.G..T.....C.....T...AT..  
 AGC AGAC..GG.A..T..A...G.T.....A.G.....CCCTTTC..GG..GGG..C...G.....A..G..  
 AGA AGAC..GG.A..T..A...G.T.....A.GT.....CC.TTTC..GG..GGG..C...G.....A..G..  
 CAG2 AGAC..GG.A..T..A...G.T.....A.GT.....CC.TTTC..GG..GGG..C...G.....A..G..  
 AGD AGAC..GG.A..T..A...G.T.....A.G.....CCCTTTC..GG..GGG..C...G.....A..G..  
 AGE AGAC..GG.A..T..A...G.T.....A.G.....CCCTTTC..GG.....CC.GCG...A..G..  
 AGI AGAC..GG.A..T..A...G.T.....A.G.....CCCT..CTC.GG..GGG..C...G.....A..G..  
 AGL AGAC..GG.A..T..A...G.T.....A.G.....CCCT..CTC.GG..GGG..C...G.....A..G..  
 AGR AG...GGCA..T.....GC.AAGT.....G.CCCTT...A..T.....TT..A...C..  
 AGS AG...GGCA..T.....GC.AAGT.....G.CCCTT...A..T.....TT..A...C..  
 Eab-A1 AGAC.AG..A..TG.....TTTGT..G.....G.....GG..GGG...AAG.G.....  
 Eab-A2 AGAC.AG..A..TG.....TTTGT..G.....G.....GG..GG...AAG.G.....  
 Eab-S7 AGACT.GG.A...G.ACC.GGT.....AA...AAAAA...GTC..TT.C..AG.....C.....T.GC....  
 68AG8 AGAC.AG..A..TGG....GG...G..AA.T.....TT...A.....TT.CA....  
 K31AG5 AGGC.AG..A..TGG....GG....AA.T.....TT...A.....TTG.ATTG..  
 16001AG3 AGAC.AG..A..TGG....GG....AA.T.....TT...A...T.....TT.ATTG..  
 TPV12MAG9 AGAC.AG..A..TGG....G.....AA..A.....TTT...A...T.....TTG..TTGC..  
 HG1HAM6 AGGC.AG..AAGTGG....G.....A.....C..TTT.....T.....AA..TT.C....  
 TE25AGBI AGAC.AG..A.....C...GG...G.....G.....G.....AA.....  
 1556AG7 AGAC.AG....T.....TA.G.....G.....G..TGGG...A.G.G.....  
 HG1PF10 AGAC.AG..A..TG.....T...T..T.....T.C.T...AGG..GGGG...AAG.G.....  
 HGIIRH165 AGAC.AG..A..TG.....T...T..T.....TG...T.....GG..GGGG...AAG.G.....  
 AG22IV AGGC.AG..A.G...AC.....A..T.....G.....G.ATGGG...AA.....  
 AG22IIIB AGGC.AG..A.G...AC.....A.....TG.....G.ATGGG...A.G.....  
 1AAG1 AGAC.AG.CAA.T.....A.GT.....TG.....G..TGG...A.....  
 Tharzianum ...C..G.C.....CC..G.....C.....C.....GGA.GGG...A..C...CA...

Fig. 1(E)

6/64

09/744502

Eab-F2 AAACC.....  
 Eab-F3 AAACC.....  
 Eab-F5 AAACC.....  
 Eab-S5 AAACC.....  
 Eab-S6 AAACC.....  
 Eab-F1 AAACC.....  
 Eab-F6 AAACC.....  
 Eab-F7 AAACC.....  
 Eab-S1 AAACC.....  
 Eab-S2 AA...T.....  
 Eab-S4 .....

521 AAA..TT..CATAT.....  
 Eab-S3 AA...TT..CATAT.....  
 RH2815 AA.C.TTTA..T.T....GGA..CC.TA.CT.CT.C.....C...TTGG..A.....  
 AGK .....  
 AGN .....  
 AGP .....  
 AGBO .....  
 AGBB .....  
 AGJ .....  
 AGG .....  
 AGC .....G...TCCGT..CTGCT..A.....  
 AGA .....G...TCCGT..CTGCT..A..TA.....  
 CAG2 .....G...TCCGT..CTGCT..A..TA.....  
 AGD .....G...TCCGT..CTGCT..A..TA.....  
 AGE .....G...TCCGT..CTGCT..A..TA.....  
 AGI .....G...TCCGT..CTGCT..A..TA.....  
 AGL .....G...TCCGT..CTGCT..A.....  
 AGR .....T..ACATA.C.A.....CC.TAA.T....A..T.....GG.AA..A..T  
 AGS .....T..ACATA.C.A.....CC.TAA.T....A..T.....GG.AA..A..T

Eab-A1 AA.C.TTTA..T.T....GGA..CC.T..T.CT..A....CT.....  
 Eab-A2 AA.C.TTTA..T.T....GGA..CC.T..T.CT..A....CT.....  
 Eab-S7 .....  
 68AG8.....TTTA..T.T....GGA..CC.....  
 K31AG5.....TTT...T.T.TA.AG.AA.C...AA.T..T..G.....G...TG....  
 16001AG3.....TTT...T.T.T.T....T...T...G.T..A.ATA..TAA..G..A.TGA..T  
 TPV12MAG9.....TTT...T.T....G.AA....A..T....A....C.AAA..GGC...AA..  
 HG1HAM6..AA...TTTA.AT.T....G...T...A.C..C..C...T..C...T.G.....  
 TE25AGBI A...TTT...T.T..AT....T...T..TG.TTCAGATAAC.AA.T.G..AATGAAGT  
 1556AG7 A..C.TTTA..T.T....GGA..C..T..CT.CT.....  
 HG1PF10 AA.C.TTTA..T.T....GGA..CC.TA.CT.CT.C.....C...TTGG.A.....  
 HGIIRH165 AA.C.TTTA..T.T....GGA..CC.T..T.CT..A....C...T.....  
 AG22IV AA.C.TTT.CAT.T.TA.....C..T..T..TT.AG.....G...TGAA.T  
 AG22IIB AA.C.TTT...TAT.T.TA....C..T..T..T..A.A.AA....T.G..AATGA..T  
 1AAG1 .....TTTA.AT.TC.....C...A..T.C..CA..T.....TTG.C..TG...T  
 Tharzianum .....AC....C.A.A..A..C..T..CT..TT...T..C...T.G...T.A.GT

Fig. 1(F)

SUBSTITUTE SHEET (RULE 26)

09/744502 2054460



09/744502

7/64

Eab-F2 .....  
 Eab-F3 .....  
 Eab-F5 .....  
 Eab-S5 .....  
 Eab-S6 .....  
 Eab-F1 .....  
 Eab-F6 .....  
 Eab-F7 .....  
 Eab-S1 .....  
 Eab-S2 .....  
 Eab-S4 .....  
  
 521 .....  
 Eab-S3 .....  
 RH2815 .....CC....T.....T.CTGTC...T.ACTTAAT.C..TATAT  
 AGK .....T.A...AAT.CCATAT..  
 AGN .....T.A...AATTC.ATAT..  
 AGP .....T.A...AATTC.ATAT..  
 AGBO .....T.A...AATTC.ATAT..  
 AGBB .....T.A...AATTC.ATAT..  
 AGJ .....T.ACTTAAT.C.....  
 AGG .....T..C..AA..CC.....  
 AGC .....T.AC.....AT..  
 AGA .....T.A.....C...AT..  
 CAG2 .....T.A.....C...AT..  
 AGD .....T.A.....C...AT..  
 AGE .....T.A.....C...AT..  
 AGI .....T.A.....C...AT..  
 AGL .....T.A.....C...AT..  
 AGR .....C.G.CACGTG.....  
 AGS .....C.G.CACGTG.....  
  
 Eab-A1 .....CCCC..T.....TG...A.CTTCTGTC...T.ACTTAATTC.ATAT..  
 Eab-A2 .....CCCC..T.....TG...A.CTTCTGTC...T.ACTTAATTC.ATAT..  
 Eab-S7 .....T.A...AAT.CCATAT..  
 68AG8 .....CC....T.C....TG.....T.C.....T.ACT....C.A.AT.T  
 K31AG5 .....C.T.GG.A.CCC...T.C....TG.T...CT.....AC.T.....C.A.AT..  
 16001AG3 AA.G..T.CATTG..AACCC...T.....T.....CT...GTC...T.ACT....C.A.A..  
 TPV12MAG9 ..TGAGTT.ATTGG.A.CCC...T.C....TG.T..CT.....AC.T..C..AATT....A..  
 HG1HAM6 .....T.C.T...A.C.....T.C.....AAT.C.....  
 TE25AGBI AA....T.C...GGGAACCC...T.C...TG.T...CT.....AC.T...T.AAT.C.....T  
 1556AG7 .....CCC...T.C....T.....CT...GTC...T.ACT....C.A.AT.T  
 HG1PF10 .....C.....T.....T.....CT...GTC...T.ACTTAAT.C...T..  
 HGIIRH165 .....CCCC..T.....TG...A.CTTCTGTC...T.ACTTAAT....T..  
 AG22IV ...GA.TT...GGGA.CCCC...TACCCCT.CTG...T.CT...AC.T..C..AA..C...T..  
 AG22IIIB .....T.....GGGA.CCCCCCT.C....TG.....T.C..T.AC.T..C..AA..C...T..  
 1AAG1 .....C.....TAC....T..T.AA.TT...T.AC...AC.....  
 Tharzianum .....CCCC..T.C....GC.G.....G..ACGT...T.A.TT....T..

Fig. 1(G)  
SUBSTITUTE SHEET (RULE 26)

FOET 20 20544260

8/64

09/744502

Eab-F2 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-F3 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-F5 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-S5 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-S6 ...ACAC.A..AA.C..CCC.A.T...CGTAT..TT.....AA..GT.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-F1 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-F6 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-F7 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-S1 ...ACAC.A..AA.C..CCC.A.T.T..GTAT..TT.....AAA..T.....T.G.AATGT.AA..TT.G.ATGT  
 Eab-S2 ...ACA.TA..AA.CT.CC..AAT.TT...A.A.T.....AAA..T.....CT.G.AATGT.AA..TT.G.ATGT  
 Eab-S4 ...ACA.TA..AA.CT.CC..AAT.TT...A.A.T.....AAA..T.....CT.G.AATGT.AA..TT.G.ATGT

521 ...ACA..A..A..CT.C...A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.A..CTT.G.ATGT  
 Eab-S3 ...ACA..A..A..CT.C...A.T.TT...A.ACT.....GCA.....CT.G.AATGT.A..CTT.G.ATGT  
 RH2815 ...A...A..A..CT.C...AAT.TT...AT..TTT.....AAAA.T.....G.AATGT.AA..T..GGATGT  
 AGK ...ACA..A..A..C..CC..A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.AA..TT.G.ATGT  
 AGN ...ACA..A..A..CT.C...A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.A..CTT.G.ATGT  
 AGP ...ACA..A..A..CT.C...A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.A..CTT.G.ATGT  
 AGB0 ...ACA..A..A..CT.C...A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.A..CTT.G.ATGT  
 AGBB ...ACA..A..A..CT.C...A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.A..CTT.G.ATGT  
 AGJ ...ACAC.A..AA.CT.C...A.T.TT...A.A.TT.....AAA.....T.T.G.AATGT.AA..TT.G.ATGT  
 AGG ...ACAC.A..AA.CT.C...A.T.T..GTAT..TT.....AAA.....CT.G.AATGT.AA..TT.G.ATGT  
 AGC .....A..AA.CT.CC..AATAT...ATA.....AA.....T.TTG.AATGT.AA..TTTG.ATGT  
 AGA .....A..AA.CT.CC..AATAT...ATA.....AA.....T...GGAATGT.AA..TTTG.ATGT  
 CAG2 .....A..AA.CT.CC..AATAT...ATA.....AA.....T...GGAATGT.AA..TTTG.ATGT  
 AGD .....A..AA.CT.CC..AATAT...ATA.....AA.....T.TTG.AATGT.AA..TTTG.ATGT  
 AGE ...A...TA.....CT.CC..AATAT...ATA.....AA.....T.TTG.AATGT.AA..TTTG.ATGT  
 AGI .....A..AA.CT.CC..A.....GTATA.T.....AAA.....T.TTG.AATGTCA...TTTG.ATGT  
 AGL .....A..AA.CT.CC..A.....GTATA.T.....AAA.....T.TTG.AATGTCA...TTTG.ATGT  
 AGR ...AC..T.....TGC..G...T.CG...T.....A...TGG...C.....G....CTT...A...  
 AGS ...AC..T.....TGC..G...T.CG...T.....A...TGG...C.....G....CTT...A...

Eab-A1 ...A...A..A..CT.C...AAT.TT...AT..TTT.....AAAA.....C..GGA.TGT.AA..T..GGATGT  
 Eab-A2 ...A...A..A..CT.C...AAT.TT...AT..TTT.....AAAA.....C..GGA.TGT.AA..T..GGATGT  
 Eab-S7 ...ACA..A..A..C..CC..A.T.TT...A.A.TT...G.AA.....CT.G.AATGT.AA..TT.G.ATGT  
 68AG8 ....CA.TA.TAAACT.C...AAT.T...TA...TTT.....AAAAATG.AA..T.G...T...AA..T..G.ATGT  
 K31AG5 ....A...TA.TAAACT.C...AAT.T...TA...TTT...AG.AAA.....T.G.AATGT.AA..T..GGATGT  
 16001AG3 CT...CA.TA.TAAACT.C...AAT.T...TA...TTT...AAAA.....T.G.AATGT.AA..T..GGATGT  
 TPV12MAG9 ...ACA.TA..AA.CT.C...AAT.T...TA...TTT...AAAA.TG.AA..T.G...T...AA..T..GGATGT  
 HG1HAM6 ...ACAC...AAAC..CC..AAT.T...TA...TTT...AAAA.....T.G.AATGT.AA..CT..GGATGT  
 TE25AGBI ....CA.TA.TAAACT.C...AAT.T...TA...TTT...AAAAATG.AA..T.G...T...AA..T..GGATGT  
 1556AG7 .TTTACAC...AAACT.C...AAT.T...TA...CTTT...AAAA.TG.AA..T.G...T...CA...TT.G.ATGT  
 HG1PF10 ....A...TA.TAAACT.C...AAT.T...TA...TTT...AAAA.TG.AA..T.G...T...AA..T..GGATGT  
 HGIIRH165 ....CA.TA.TAAACT.C...AAT.T...TA...TTT...AAAA.....C..GGA.TGT.AA..T..GGATGT  
 AG22IV CT..A.A.TAC.AAA...CCC.AAT.T...TA...TTT...AAAA.TG.AA..T.G...T...AA..T..GGATGT  
 AG22IIIB CT..A.A.TA.TAAA...CCC.AAT.T...TA...TTT...AAAA.....T.G.AATGT.AA..T..GGATGT  
 1AAG1 ...ACACT.CTA.....C.....T...TA.A.TT...AAA.....CT.G.AATGT.AA..TT.G.ATGT  
 Tharzianum CTT.ACA.....GC.....T...C.T.....GAGCAAAAA.....TT.....C.....

Fig. 1(H)

SUBSTITUTE SHEET (RULE 26)

09744502 PCT/GB99/02406

09/744502

9/64

Eab-F2	AACG..CAT.C.A.TTA.GG.AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-F3	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-F5	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S5	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S6	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-F1	AACG..CAT.C.A.TTAA...AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-F6	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-F7	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S1	AACG..CAT.C.A.TTA.G..AACT	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S2	AAC..ACAT.CT..TTA...AACT.	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S4	AAC..ACAT.CT..TTA...AACT.	end of ITS1	AAG.TTTCACAACGGATCTCTGGCTCTCGC

521	AACG..CAT.C.A.TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S3	AACG..CAT.C.A.TTA...CA.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
RH2815	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGK	AACG..CAT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGN	AACG..CAT.C.A.TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGP	AACG..CAT.C.A.TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGBO	AACG..CAT.C.A.TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGBB	AACG..CAT.C.A.TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGJ	AACG..CAT.C.A.TTA.G..AACT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGG	AACG..CAT.C.A.TTA.G..AACT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGC	AACG..CAT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGA	AACGG..AT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
CAG2	AACGG..AT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGD	AACG..CAT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGE	AACG..CAT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGI	AAC..ACAT.CTA.T.AA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGL	AAC..ACAT.CTA.T.AA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGR	A...A.A..CTAA..AA...A.C.	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AGS	AA...AC.T...AA..AA...A.C.	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC

Eab-A1	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTGCAACAACGGATCTCTGGCTCTCGC
Eab-A2	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Eab-S7	AACG..CAT.CT..TTAA...A.CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
68AG8	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
K31AG5	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
16001AG3	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
TPV12MAG9	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
HG1HAM6	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
TE25AGBI	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
1556AG7	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
HG1PF10	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
HG1IRH165	AAC..ACAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AG22IV	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
AG22IIB	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
1AAG1	AACG..CAT.CTAAT.A.....CT	[end of ITS1]	AAG.TTTCACAACGGATCTCTGGCTCTCGC
Tharzianum	AA...A.ATG..AAT.....CAA..	[end of ITS1]	AA.CTTTCAACAACGGATCTCTGGTCTGGC

Fig. 1(I)

SUBSTITUTE SHEET (RULE 26)

09744502-071301

09/744502

10/64

F2 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
F3 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
F5 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S5 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S6 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
F1 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
F6 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
F7 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S1 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
521 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S3 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S7 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S2 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
S4 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
A2 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
A1 ATCGATGAAA AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
RH2815 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGK ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGN ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGP ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGB0 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGBB ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGJ ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGG ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGC ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGA ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
CAG2 ATCGATGAAG AACGCAGCGA AACCCGATAA GTAATGTGAA TTGCAGAATT  
AGD ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGE ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGI ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGL ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGR ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AGS ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
68AG8 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
K31AG5 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
16001AG3 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
TPV12MAG9 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
HG1HAM6 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
TE25AGBI ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
1556AG7 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
HG1PF10 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
HGIIRH165 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AG22IV ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
AG22IIB ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
1AAG1 ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT  
THARZIANU ATCGATGAAG AACGCAGCGA AATGCGATAA GTAATGTGAA TTGCAGAATT

Fig. 1(J)

SUBSTITUTE SHEET (RULE 26)

09744502-100

09/744502

11/64

F2 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 F3 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 F5 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S5 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S6 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 F1 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 F6 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 F7 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S1 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 521 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S3 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S7 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S2 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 S4 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 A2 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 A1 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 RH2815 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGK CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGN CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGP CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGBO CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGBB CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGJ CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGG CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGC CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGA CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 CAG2 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGD CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGE CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGI CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGL CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGR CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AGS CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 68AG8 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 K31AG5 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 16001AG3 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 TPV12MAG9 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 HG1HAM6 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 TE25AGBI CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 1556AG7 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 HG1PF10 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 HGIIRH165 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AG22IV CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 AG22IIB CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT  
 1AAG1 CAGTGAATCA TCGAATCTTT GAACGCACCT TCGCTCC.TT ...GGTATTCCTT..  
 THARZIANU CAGTGAATCA TCGAATCTTT GAACGCACAT TCGC.CCG.. CCAG.TATTC.T.GG

Fig. 1(K)

SUBSTITUTE SHEET (RULE 26)

09744502.0130

09/744502

12/64

F2 GGAGCATGCC TGT TT..GAG..TA TCAT  
F3 GGAGCATGCC TGT TT..GAG..TA TCAT  
F5 GGAGCATGCC TGT TT..GAG..TA TCAT  
S5 GGAGCATGCC TGT TT..GAG..TA TCAT  
S6 GGAGCATGCC TGT TT..GAG..TA TCAT  
F1 GGAGCATGCC TGT TT..GAG..TA TCAT  
F6 GGAGCATGCC TGT TT..GAG..TA TCAT  
F7 GGAGCATGCC TGT TT..GAG..TA TCAT  
S1 GGAGCATGCC TGT TT..GAG..TA TCAT  
521 GGAGCATGCC TGT TT..GAG..TA TCAT  
S3 GGAGCATGCC TGT TT..GAG..TA TCAT  
S7 GGAGCATGCC TGT TT..GAG..TA TCAT  
S2 GGAGCATGCC TGT TT..GAG..TA TCAT  
S4 GGAGCATGCC TGT TT..GAG..TA TCAT  
A2 GGAGCATGCC TGT TT..GAG..TA TCAT  
A1 GGAGCATGCC TGT TT..GAG..TA TCAT  
RH2815 GGAGCATGCC TGT TT..GAG..TA TCAT  
AGK GGAGCATGCC TGT TT..GAG..TA TCAT  
AGN GGAGCATGCC TGT TT..GAG..TA TCAT  
AGP GGAGCATGCC TGT TT..GAG..TA TCAT  
AGBO GGAGCATGCC TGT TT..GAG..TA TCAT  
AGBB GGAGCATGCC TGT TT..GAG..TA TCAT  
AGJ GGAGCATGCC TGT TT..GAG..TA TCAT  
AGG GGAGCATGCC TGT TT..GAG..TA TCAT  
AGC GGAGCATGCC TGT TT..GAG..TA TCAT  
AGA GGAGCATGCC TGT TT..GAG..TA TCAT  
CAG2 GGAGCATGCC TGT TT..GAG..TA TCAT  
AGD GGAGCATGCC TGT TT..GAG..TA TCAT  
AGE GGAGCATGCC TGT TT..GAG..TA TCAT  
AGI GGAGCATGCC TGT TT..GAG..TA TCAT  
AGL GGAGCATGCC TGT TT..GAG..TA TCAT  
AGR GGAGCATGCC TGT TT..GAG..TA TCAT  
AGS GGAGCATGCC TGT TT..GAG..TA TCAT  
68AG8 GGAGCATGCC TGT TT..GAG..TA TCAT  
K31AG5 GGAGCATGCC TGT TT..GAG..TA TCAT  
16001AG3 GGAGCATGCC TGT TT..GAG..TA TCAT  
TPV12MAG9 GGAGCATGCC TGT TT..GAG..TA TCAT  
HG1HAM6 GGAGCATGCC TGT TT..GAG..TA TCAT  
TE25AGBI GGAGCATGCC TGT TT..GAG..TA TCAT  
1556AG7 GGAGCATGCC TGT TT..GAG..TA TCAT  
HG1PF10 GGAGCATGCC TGT TT..GAG..TA TCAT  
HGI1RH165 GGAGCATGCC TGT TT..GAG..TA TCAT  
AG22IV GGAGCATGCC TGT TT..GAG..TA TCAT  
AG22IIB GGAGCATGCC TGT TT..GAG..TA TCAT  
1AAG1 GGAGCATGCC TGT TT..GAG..TA TCAT  
Tharzianum CGGGCATGCC TGT..CCGAGCG.. TCAT

Fig. 1(L)

SUBSTITUTE SHEET (RULE 26)

09744502.07301

09/744502

13/64




Fig. 2

14/64




Fig. 3



09/744502

15/64




Fig. 4

16/64




Fig. 5

09/744502

17/64




Fig. 6

09/744502

18/64




Fig. 7

09/744502

19/64




Fig. 8

09/744502

20/64




Fig. 9

09/744502

21/64




Fig. 10

09/744502

22/64

	0 Days	3 Days	5Days
Me 8.2 (+)	3	3	3
RAIb (+)	1	2	3
Fusarium (+)	1	3	3
C (-)	0,1	0,1	0,1
Eab-F5/Me 8.2	0,1	0,1	0,1
Eab-F5/Me 8.2	0,1	0,1	1
Eab-S6/Me 8.2	0,1	0,1	0,1
Eab-S6/Me 8.2	0,1	0,1	0,1
Eab-F5/RAIb Si	0,1	2	3
Eab-F5/RAIb C	0,1	0,1	1
Eab-S6/RAIb Si	0,1	0,1	0,1
Eab-S6/RAIb C	0,1	0,1	0,1
Eab-F5/Fusari	1	2	3
Eab-F5/Fusari	0,1	0,1	0,1
Eab-S6/Fusariu	0,1	0,1	0,1
Eab-S6/Fusariu	0,1	0,1	0,1




Fig. 11



09/744502

23/64

	0 days	4 days	7 days
Me 8.2 (+)	1	3	5
RAlb (+)	2	5	5
Fusarium (+)	1	2	3
C(-)	0,1	0,1	0,1
Eab-S1/Me 8	0,1	1	2
Eab-S1/Me 8	0,1	0,1	0,1
Eab-S7/Me 8	2	3	4
Eab-S7/Me 8	1	1	1
Eab-S1/Ralb	1	3	4
Eab-S1/Ralb	0,1	0	0,1
Eab-S7/Ralb	2	4	4
Eab-S7/Ralb	0,1	0,1	0,1
Eab-S1/Fusa	0,1	0,1	1
Eab-S1/Fusa	0,1	0,1	0,1
Eab-S7/Fusa	1	1	2
Eab-S7/Fusa	0,1	0,1	0,1




Fig. 12

SUBSTITUTE SHEET (RULE 26)

09/744502

24/64

	0 Days	3 Days	6 Days
Me 8.2 (+)	4	4	5
RAIb (+)	3	4	4
Fusarium (+)	1	2	3
C (-)	0,1	0,1	0,1
Eab-F1/Me 8.2	4	5	5
Eab-F1/Me 8.2	0,1	1	1
Eab-F7/Me 8.2	3	4	5
Eab-F7/Me 8.2	0,1	0,1	0,1
Eab-F1/RAIb Si	5	5	5
Eab-F1/RAIb C	0,1	0,1	0,1
Eab-F7/RAIb Si	5	5	5
Eab-F7/RAIb C	0,1	0,1	0,1
Eab-F1/Fusari	1	3	3
Eab-F1/Fusari	0,1	0,1	0,1
Eab-F7/Fusari	2	3	3
Eab-F7/Fusari	1	1	1

## SUGAR BEET PROTECTION TEST




Fig. 13

25/64

## PROTECTION TEST 1

Negative control without fungus (C-), positive control with pathogenic strain Me 8.2, and two protective binucleate strains, F2 and  $\alpha$ B, on tomato seedlings.




Fig. 14

SUBSTITUTE SHEET (RULE 26)

09744502-07301

26/64

## PROTECTION TEST 1

Protection of binucleate strains F2 (up) and  $\alpha$ B (down) from pathogenic strain Me 8.2 on tomato seedlings when applied consecutively (right) but not when applied simultaneously (left).




Fig. 15

27/64

## PROTECTION TEST 3

(Up) Negative control without fungus (C-), positive control with pathogenic strain Me 8.2. (Down) Negative control without fungus (C-), positive control with pathogenic strain pin.




Fig. 16

SUBSTITUTE SHEET (RULE 26)

FOET 20 205774502



28/64

## PROTECTION TEST 3

Protection of binucleate strain **Eab-F2** against pathogenic **Rhizoctonia Me 8.2** (up) and **pin** (down) strains on pine seedlings when applied consecutively (right) and simultaneously (left).




Fig. 17

SUBSTITUTE SHEET (RULE 26)

09/744502-01301

29/64

## PROTECTION TEST 3

Protection of binucleate strain **Eab-aB** against pathogenic **Rhizoctonia** Me 8.2 (up) and **pin** (down) strains on pine seedlings when applied consecutively (right) and simultaneously (left).




Fig. 18

30/64

## PROTECTION TEST 3

(Up) Negative control without fungus (C-), positive control with pathogenic strain **pin** (Down) Negative control without fungus (C-), positive control with pathogenic strain **Me 8.2**.





Fig. 19



## PROTECTION TEST 3

(Up) Protection of binucleate strain **Eab-aB** against pathogenic **Me 8.2** strain on pine seedlings when applied consecutively (left) and simultaneously (right). (Down) Protection of binucleate strain **Eab-aB** against pathogenic **pin** strain on pine seedlings when applied consecutively (left) and simultaneously (right).





Fig. 20

32/64

## PROTECTION TEST 3

(Up) Protection of binucleate strain **Eab-F2** against pathogenic **Me 8.2** strain on pine seedlings when applied consecutively (left) and simultaneously (right). (Down) Protection of binucleate strain **Eab-F2** against pathogenic **pin** strain on pine seedlings when applied consecutively (left) and simultaneously (right).





Fig. 21

PROTECTION TEST 3 - PINE SEEDLINGS




Fig. 22

34/64

FOET 20" 20544260

PROTECTION TEST 3 - PINE SEEDLINGS




Fig. 23

35/64

PROTECTION TEST 3 - PINE SEEDLINGS




Fig. 24



36/64

## PROTECTION TEST 4

Protection of binucleate strains **F2** (up) and **S7** (down) from pathogenic *Fusarium* strain on wheat seedlings when applied consecutively (left) and simultaneously (right).




Fig. 25

SUBSTITUTE SHEET (RULE 26)

09/744502-01301

37/64

## PROTECTION TEST 4

Protection of binucleate strains **S6** (up) and **F5** (down) from pathogenic **Rhizoctonia RAG2** strain on wheat seedlings when applied consecutively (left) and simultaneously (right).



Fig. 26

38/64

## PROTECTION TEST 4

(up) Negative control without fungus (C-), positive control with pathogenic *Fusarium* strain

(down) Negative control without fungus (C-), positive control with pathogenic *Rhizoctonia* RAG2 strain




Fig. 27



39/64

## PROTECTION TEST 4

General view of experimental greenhouse design (up)  
and detail of early symptoms caused by pathogenic  
strain *Fusarium sp.* (Down)



Fig. 28

40/64

## PROTECTION TEST 4

(up) Negative control without fungus (C-), positive control with pathogenic *Fusarium* strain

(down) Negative control without fungus (C-), positive control with pathogenic *Rhizoctonia* RAG2 strain



Fig. 29

41/64

## PROTECTION TEST 4

Protection of binucleate strain **F5** from pathogenic *Fusarium* *sp.* strain on wheat plants when applied consecutively (up) and simultaneously (down).



Fig. 30

SUBSTITUTE SHEET (RULE 26)

09/744502 20544260

42/64

## PROTECTION TEST 4

Protection of binucleate strain **S6** from pathogenic *Fusarium* *sp.* strain on wheat plants when applied consecutively (up) and simultaneously (down).



Fig. 31

SUBSTITUTE SHEET (RULE 26)

T06T20 20544/60

43/64

## PROTECTION TEST 5

(up) Negative control without fungus (C-), positive control with pathogenic *Rhizoctonia* RAG2 strain

(down) Protection of binucleate strains F3 from pathogenic *Rhizoctonia* RAG2 strain on rape seedlings when applied consecutively (left) and simultaneously (right).




Fig. 32

SUBSTITUTE SHEET (RULE 26)

09/744502-01301



44/64

## PROTECTION TEST 5

Protection of binucleate strains **S6** (up) and **F6** (down) from pathogenic *Rhizoctonia* **RAG2** strain on rape seedlings when applied consecutively (left) and simultaneously (right).




Fig. 33

45/64

## PROTECTION TEST 5

Protection of binucleate strains **S1** (up) and **S7** (down) from pathogenic **Rhizoctonia RAG2** strain on rape seedlings when applied consecutively (left) and simultaneously (right).




Fig. 34

SUBSTITUTE SHEET (RULE 26)

FOET 0 2054460

46/64

## PROTECTION TEST 6

(up) Negative control without fungus (C-), positive control with pathogenic *Fusarium* strain

(down) Negative control without fungus (C-), positive control with pathogenic *Rhizoctonia* RAG2 strain




Fig. 35



47/64

## PROTECTION TEST 6

Negative control without fungus (C-), positive control with pathogenic *Rhizoctonia meloni* 8.2 strain




Fig. 36

48/64

## PROTECTION TEST 6

Protection of binucleate strains **S1** (up) and **S7** (down) from pathogenic *Rhizoctonia* **RAG2** strain on alfalfa seedlings when applied consecutively (right) and simultaneously (left).




Fig. 37

SUBSTITUTE SHEET (RULE 26)

09/744502-071301

49/64

## PROTECTION TEST 6

Protection of binucleate strains S1 (up) and S7 (down) from pathogenic *Rhizoctonia meloni* 8.2 strain on alfalfa seedlings when applied consecutively (right) and simultaneously (left).




Fig. 38

SUBSTITUTE SHEET (RULE 26)

09/744502-01301

50/64

## PROTECTION TEST 6

Protection of binucleate strains **S1** (up) and **S7** (down) from pathogenic *Fusarium* strain on alfalfa seedlings when applied consecutively (right) and simultaneously (left).




Fig. 39



51/64

## PROTECTION TEST 7

Negative control with non-sterilized garlic bulbs  
(C-), infected with pathogenic *Penicillium* sp.  
strain.




Fig. 40

52/64

## PROTECTION TEST 7

Protection of binucleate strains **F1** (up) and **S4** (down) from pathogenic *Penicillium* strain on garlic bulbs when applied directly on the surface of the bulb




Fig. 41

53/64

## PROTECTION TEST 7

Protection of binucleate strains **F7** (up) and **S6** (down) from pathogenic *Penicillium* strain on garlic bulbs when applied directly on the surface of the bulb




Fig. 42



54/64

## PROTECTION TEST 8

Protection of binucleate strains **Eab-F1** (up) and **Eab-F2** (down) from pathogenic *Fusarium sp.* strain on sugar beet seedlings when applied consecutively (right) and simultaneously (left).




Fig. 43



55/64

## PROTECTION TEST 8

Negative control without fungus (C-), positive control with pathogenic *Fusarium sp.* strain




Fig. 44

56/64

## PROTECTION TEST 8

Protection of binucleate strains **Eab-F1** (up) and **Eab-F3** (down) from pathogenic **Rhizoctonia RAG2** strain on sugar beet seedlings when applied consecutively (right) and simultaneously (left).




Fig. 45

57/64

## PROTECTION TEST 8

Protection of binucleate strains **Eab-F1** (up) and **Eab-S7** (down) from pathogenic **Rhizoctonia meloni** 8.2 strain on sugar beet seedlings when applied consecutively (right) and simultaneously (left).




Fig. 46



58/64

## PROTECTION TEST 8

Protection of binucleate strains **Eab-S1** (up) and **Eab-S7** (down) from pathogenic **Rhizoctonia RAlb** strain on sugar beet seedlings when applied consecutively (right) and simultaneously (left).




Fig. 47

SUBSTITUTE SHEET (RULE 26)

09/744502-031

59/64

## PROTECTION TEST 8

(Up) Negative control without fungus (C-), positive control with pathogenic strain Me 8.2. (Down) Negative control without fungus (C-), positive control with pathogenic strain *Rhizoctonia* RAlb.




Fig. 48

SUBSTITUTE SHEET (RULE 26)

60/64

## PROTECTION TEST 9

(Up) Negative control without fungus (C-), positive control with pathogenic strain **RA1b**. (Down) Negative control without fungus (C-), positive control with pathogenic strain *Fusarium sp.*




Fig. 49



61/64

## PROTECTION TEST 9

Negative control without fungus (C-), positive control with pathogenic strain Me 8.2.



Fig. 50

62/64

## PROTECTION TEST 9

Protection of binucleate strains **Eab-F1** (up) and **Eab-F6** (down) from pathogenic **Rhizoctonia** RAlb strain on onion seedlings when applied consecutively (right) and simultaneously (left).




Fig. 51



63/64

## PROTECTION TEST 9

Protection of binucleate strains **Eab-F3** (up) and **Eab-F6** (down) from pathogenic *Rhizoctonia* Me 8.2 strain on onion seedlings when applied consecutively (right) and simultaneously (left).




Fig. 52

64/64

## PROTECTION TEST 9

Protection of binucleate strains Eab-F3 (up) and Eab-F6 (down) from pathogenic *Fusarium* sp. strain on onion seedlings when applied consecutively (right) and simultaneously (left).




Fig. 53

USA

DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: Protection of Plants Using Rhizoctonia

which is described and claimed in:

[ ] the attached specification

[X] the specification in application Serial No.

09/744,502

filed 1/24/01

[x] PCT International Application No.

PCT/GB99/02406

filed July 23, 1999

(if applicable) and amended on

under Article 19 PCT

and on

under Article 34 PCT

I hereby state that I have reviewed and understand the contents of the above-identified application specification, including the claims, as amended by any amendment specifically referred to herein.

I acknowledge the duty to disclose all information known to me that is material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code §119 of any foreign applications(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Number	Country	Date Filed	Priority Claimed
<u>9816265.4</u>	<u>United Kingdom</u>	<u>July 24, 1998</u>	[x]yes [ ]no
<u>                    </u>	<u>                    </u>	<u>                    </u>	[ ]yes [ ]no
<u>                    </u>	<u>                    </u>	<u>                    </u>	[ ]yes [ ]no
<u>                    </u>	<u>                    </u>	<u>                    </u>	[ ]yes [ ]no

I hereby claim the benefit under Title 35, United States Code §1.20, of the United States Application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information that is material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56, and which became available to me between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status (patented, pending, abandoned)
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment or both under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardize the validity of the application or any patent issued thereon.

09/744,502 "0" 2005/04/26

## I hereby appoint

ALTHER, Robert F.	31,810	HONG, Patricia E.	34,373	PATEL, Binal J.	42,065
BANNER, Donald W.	17,037	HOSCHEIT, Dale H.	19,090	PATHAK, Ajay S.	38,266
BANNER, Mark T.	29,888	IWANICKI, John P.	34,628	PAYNE, Stephen S.	35,316
BANNER, Pamela I.	33,644	JACKSON, Thomas H.	29,808	PETERSON, Thomas L.	30,969
BECKETT, William W.	18,262	KAGAN, Sarah A.	32,141	POTENZA, Joseph M.	28,175
BODNER, Jordan	42,338	KATZ, Robert S.	36,402	PRATT, Thomas K.	37,210
BUROW, Scott A.	42,373	KLEIN, William J.	43,719	RENK, Christopher J.	33,761
CALLAHAN, James V.	20,095	KRAUSE, Joseph P.	32,578	RESIS, Robert H.	32,168
CHANG, Steve S.	42,402	LINEK, Ernest V.	29,822	RIVARD, Paul M.	43,446
COHAN, Gregory J.	40,959	MAGOON, Sumeet	43,769	SCHAD, Steve P.	32,550
COOPERMAN, Marc S.	34,143	MALONE, Dale A.	32,155	SHIFLEY, Charles W.	28,042
CURTIN, Joseph P.	34,571	MANNAVA, Ashok K.	P45,301	SKERPON, Joseph M.	29,864
DAWSON, John R.	39,504	McDERMOTT, Peter D.	29,411	SPAR, Elizabeth	45,123
DEMOOR, Laura J.	39,654	McKEE, Christopher L.	32,384	STOCKLEY, D. J.	34,257
EVANS, Thomas L.	35,805	McKIE, Edward F.	17,335	VAN ES, J. Pieter	37,746
FEDOROCKO, Gary D.	35,509	MEDLOCK, Nina L.	29,673	WILLIAMS, Kathleen M.	34,380
FISHER, Daniel E.	34,162	MEECE, Timothy C.	38,553	WITCOFF, Sheldon W.	17,399
FISHER, William J.	32,133	MEEKER, Frederic M.	35,282	WOLFF, Kevin A.	42,233
GLEBOCKI, Christopher R.	38,800	MILLER, Charles L.	43,805	WOLFFE, Franklin D.	19,724
HANLON, Brian E.	40,449	MITRIUS, Janice V.	43,808	WOLFFE, Susan A.	33,568
HART, Robert P.	35,184	MORENO, Christopher P.	38,566	WRIGHT, Bradley C.	38,061
HEMMENDINGER, Lisa M.	42,653	NELSON, Jon O.	24,566		
HILLMAN, Lisa	43,673	NIEGOWSKI, James A.	28,331		

jointly, and each of them severally, my attorneys and attorney, with full power of substitution, delegation and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent and to transact all business in the Patent and Trademark Office connected therewith. Please direct all correspondence and telephone calls to:

Ernest V. Linek  
Banner & Witcoff, Ltd.  
28 State Street, 28th Floor  
Boston, MA 02109

Customer Number: 22910  
Tel: (617) 227-7111  
Fax: (617) 227-4399

1 - 00  
Full Name of first Inventor:

Residence:

Citizenship:

Post Office Address:

Rubio Susan, Victor

Madrid, 28008, Spain ESX

Spanish

Cuesta San Vicente 20,

Madrid, 28008,

Spain

2 - 00  
Full Name of second Inventor:

Residence:

Citizenship:

Post Office Address:

Salazar Torres, Oscar

Madrid, 28047, Spain ESX

Spanish

Valmojado 233, 1A, Madrid, 28047,

Spain

3 - 00  
Full Name of third Inventor:

Residence:

Citizenship:

Post Office Address:

Julian Esquivias, Maria

Alcobendas, 28100, Spain ESX

Spanish

Paseo de la Chopera, 2B,

Alcobendas, 28100,

Spain

4 - 00  
Full Name of fourth Inventor:

Residence:

Citizenship:

Post Office Address:

Gonzalez Garcia, Vicente

Madrid, 28022, ESX

Spanish

Colonia Occidente 7,

Madrid, 28022,

Spain

5 - 00  
Full Name of fifth Inventor:  
Residence:  
Citizenship:  
Post Office Address:

Munoz Gomez, Ramona  
Casas Ibanez, 02200, Spain ESX  
Spanish  
Medico 10,  
Casas Ibanez, 02200, Spain

6 - 00  
Full Name of sixth Inventor:  
Residence:  
Citizenship:  
Post Office Address:

Gomez-Acebo Gullon, Eduardo  
Pozuelo, 28224, Spain ESX  
Spanish  
Sanchidrian 18, Pozuelo, 28224,  
Spain

7 - 00  
Full Name of seventh Inventor:  
Residence:  
Citizenship:  
Post Office Address:

Lopez Corcoles, Horacio  
Albacete, 02006, Spain ESX  
Spanish  
Historia 31, Albacete, 02006,  
Spain

Signature of first inventor:



Date:

March, 14th, 2001

Signature of second inventor:



Date:

March, 14th, 2001

Signature of third inventor:



Date:

March, 14th, 2001

Signature of fourth inventor:



Date:

March, 14th, 2001


Signature of fifth inventor:



Date:

March, 14th, 2001


Signature of sixth inventor:



Date:

March, 14th, 2001

Signature of seventh inventor:



Date:

March, 14th, 2001

PCT Rec'd 13 JUL 2001

Docket No. 4512/00012

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

**APPLICANT:** Rubio Susan et al. **EXAMINER:** Unknown  
**SERIAL NO.:** 09/744,502 **GROUP:** Unknown  
**FILED:** 24 January 2000  
**FOR:** Protection of Plants Using Rhizoctonia

ASSISTANT COMMISSIONER FOR PATENTS  
WASHINGTON, DC 20231  
BOX PCT

Sir:


## STATEMENT

In response to Notification to Comply with Requirement for Patent Applications, I declare that the contents of the paper, the computer readable form and the compact disc attached are the same and do not contain any new matter in connection with the above-identified application.

I hereby declare that all statements made herein of my own knowledge are true and that all statements were made with the knowledge that willful false statement and the like so made are punishable by fine or imprisonment or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Respectfully submitted,

MARKS &amp; CLERK

  
Name: **MR. D. ICELY**  
Title: **TECHNICAL ASSISTANT**

Date: 6/7/01

## SEQUENCE LISTING

<110> Rubio Susan, Victor  
Salazar Torres, Oscar  
Julain Esquivivias, Maria  
Gonzales Garcia, Vincente  
Munoz Gomez, Ramona  
Gomez-Acebo Gullon, Eduardo  
Lopez Corcoles, Horacio

<120> Protection of Plants Using Rhizoctonia

<130> USP79179

<140> US 09/744,502

<141> 2001-01-24

<150> PCT/GB99/02406

<151> 1999-07-23

<160> 28

<170> PatentIn version 3.0

<210> 1

<211> 15

<212> DNA

<213> Rhizoctonia sp.

<400> 1

ccctattaag gggca

15

<210> 2

<211> 15

<212> DNA

<213> Rhizoctonia sp.

<400> 2

ccgtaaaaaa gtctt

15

<210> 3

<211> 20

<212> DNA

<213> Rhizoctonia sp.

<400> 3  
ccacacaaac ccattgtatt 20

<210> 4  
<211> 18  
<212> DNA  
<213> Rhizoctonia sp.

<400> 4  
aattgttcct ttgggcac 18

<210> 5  
<211> 16  
<212> DNA  
<213> Rhizoctonia sp.

<400> 5  
ggaaccttta ttggac 16

<210> 6  
<211> 20  
<212> DNA  
<213> Rhizoctonia sp.

<400> 6  
ctcaatttat tttaaaacga 20

<210> 7  
<211> 19  
<212> DNA  
<213> Rhizoctonia sp.

<400> 7  
ttgaatgaat atagagttg 19

<210> 8  
<211> 21  
<212> DNA  
<213> Rhizoctonia sp.



<400> 8  
tcgaatgaac aatggggatc g 21

<210> 9  
<211> 11  
<212> DNA  
<213> Rhizoctonia sp.

<400> 9  
tcctccggga g 11

<210> 10  
<211> 13  
<212> DNA  
<213> Rhizoctonia sp.

<400> 10  
ttcttcattg agg 13

<210> 11  
<211> 20  
<212> DNA  
<213> Rhizoctonia sp.

<400> 11  
agacggagga ccgtaaaaaa 20

<210> 12  
<211> 16  
<212> DNA  
<213> Rhizoctonia sp.

<400> 12  
ttccgtctat taaacc 16

<210> 13  
<211> 26  
<212> DNA  
<213> Rhizoctonia sp.

<400> 13

acacaaaccc cattgtatatt aaattg 26

<210> 14

<211> 26

<212> DNA

<213> Rhizoctonia sp.

<400> 14

acataaactc caatttaata aatctg 26

<210> 15

<211> 242

<212> DNA

<213> Rhizoctonia sp.

<400> 15

agaggaagta aaagtcgtaa caaggtttcc gtaggtgaac ctgcggaagg atcattattg 60

aatgaatata gagttggttg tcgctggctc ctccgggagc atgtgcacgc tttctctttc 120

atccacacac acctgtgcac ttgtgagacg gaggaccgta aaaaagtctt ccgtctatta 180

aaccacacaa accccattgt atttaaattg aatgtaattg atgtaacgca tcattaggaa 240

ct 242

<210> 16

<211> 243

<212> DNA

<213> Rhizoctonia sp.

<400> 16

ttagaggaag taaaagtcgt aacaaggttt ccgtaggtga acctgcggaa ggatcattat 60

tgaatgaata tagagttggt tgtecgctggc tcctccggga gcatgtgcac gctttctctt 120

tcacccacac acacctgtgc acttgtgaga cggaggaccg taaaaaagtc ttccgtctat 180

taaaccacac aaacccatt gtatttaaatt tgaatgtaat tgatgtaacg catcattaga 240

act 243

<210> 17  
 <211> 238  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 17  
 ggaagtaaaa gtcgtaacaa ggtttccgta ggtgaacctg cggaaggatc attattgaat 60  
 gaatatagag ttggttgctg ctggctcctc cgggagcatg tgcacgcttt ctctttcatc 120  
 cacacacacc tgtgcacttg tgagacggag gaccgtaaaa atgtcttccg tctattaaac 180  
 cacacaaacc ccattgtatt taaattgaat gtaattgatg taacgcatca ttagaact 238

<210> 18  
 <211> 236  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 18  
 tggataaagt cgtaacaagg tttccgtagg tgaacctgcg gaaggatcat tattgaatga 60  
 atatatagagtt ggttgctgct ggctcctccg ggagcatgtg cacgctttct ctttcatcca 120  
 cacacacctg tgcacttggt agacggagga ccgtaaaaaa gtcttccgtc tattaaacca 180  
 cacaaacccc attgtattta aattgaatgt aattgatgta acgcatcatt agaact 236

<210> 19  
 <211> 237  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 19  
 gggataaaaag tcgtaacaag gtttccgtag gtgaacctgc ggaaggatca ttattgaatg 60  
 aatatagagt ttggttgctg ttggctcctcc gggagcatgt gcacgctttc tctttcatcc 120  
 acacacacgt gtgcacttgt gagacggagg accgtaaaaa agtcttccgt ctattaaacc 180  
 acacaaaccc catcgtattt aagttagatg taattgatgt aacgcatcat tagaact 237

<210> 20

&lt;211&gt; 245

&lt;212&gt; DNA

&lt;213&gt; Rhizoctonia sp.

&lt;400&gt; 20

ttagagggga ataaaaagtc gtaacaaggt ttccgtaggt gaacctgcgg aaggatcatt 60

attgaatgaa tatagagttg gttgtcgctg gtcctccgg gagcatgtgc acgctttctc 120

tttcatccac acacacctgt gcacttgtga gacggaggac cgtaaaaaag tcttccgtct 180

attaaaccac acaaacccca ttgtatttaa attgaatgta attgatgtaa cgcattcatta 240

aaact 245

&lt;210&gt; 21

&lt;211&gt; 246

&lt;212&gt; DNA

&lt;213&gt; Rhizoctonia sp.

&lt;400&gt; 21

ttagagggga agtaaaaagt cgtaacaagg ttccgtagg tgaacctgcg gaaggatcat 60

ttattgaatga atatagagtt ggttgtcgct ggctcctccg ggagcatgtg cacactttct 120

ctttcatcca cacacacctg tgcacttgtg agacggagga ccgtaaaaaa gtcttccgtc 180

tattaaacca cacaaacccc attgtattta aattgaatgt aattgatgta acgcattcatt 240

agaact 246

&lt;210&gt; 22

&lt;211&gt; 252

&lt;212&gt; DNA

&lt;213&gt; Rhizoctonia sp.

&lt;400&gt; 22

ttcccttttt tagaggaagt aaaagtcgta acaaggtttc cgtaggtgaa cctgcggaag 60

gatcattatt gaatgaatat agagttgggt gtcgctgggt cctccgggag catgtgcacg 120

ctttctcttt catccacaca cacctgtgca cttgtgagac ggaggaccgt aaaaaagtct 180

tccgtctatt aaaccacaca aaccccattg tatttaaatt gaatgtaatt gatgtaacgc 240

atcattagaa ct 252

<210> 23  
 <211> 230  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 23  
 agtcgtaaca aggtttccgt aggtgaacct gcggaaggat cattattgaa tgaatataga 60  
 gttggttgtc gctggctcct cgcgggagca tgtgcacgct ttctctttca tccacacaca 120  
 cctgtgcact tgtgagacgg aggaccgtaa aaaagtcttc cgtctattaa accacacaaa 180  
 cccattgta tttaaattga atgtaattga tgtaacgcat cattagaact 230

<210> 24  
 <211> 232  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 24  
 gtcgtaacaa gggtttccgta ggtgaacctg cggaaggatc attatcgaat gaacaatggg 60  
 gatcggttgt cgctggcttc ttcattgagg catgtgcacg cctttctcta tttcatccac 120  
 acacacctgt gaacttgtga gacggatagt agtaaaaaaa gtcttcgtct gtcaatacat 180  
 aaactccaat ttaataaatc tgaatgtaat tgatgtaaca catctttaa ct 232

<210> 25  
 <211> 226  
 <212> DNA  
 <213> Rhizoctonia sp.

<400> 25  
 gtcgtaacaa gggtttccgta ggtgaacctg cggaaggatc attatcgaat gaacaatggg 60  
 gatcggttgt cgctggcttc ttcattgagg catgtgcacg cctttctcta tttcatccac 120  
 acacacctgt gaacttgtga gacggaggac cgtaaaaaag tcttctgtct acataaactc 180

càatttāata aatctgaatg taattgatgt aacacatctt taaact 226

<210> 26

<211> 173

<212> DNA

<213> Rhizoctonia sp.

<400> 26

agagtcgggt gtcgctgggt gtctttggca gcatgtgcac gccttctctt ttcattccaca 60

caccctgtg cacttgtgag actggaggcc gtaaaaagcc ttcagtctgc taaattcata 120

tacaaactca tttaattgaa ctgaatgtac ttgatgtaac gcatcattaa act 173

<210> 27

<211> 230

<212> DNA

<213> Rhizoctonia sp.

<400> 27

aaagtcgtaa caaggtttcc gtaggtgaac ctgcggaagg atcattattg aatgaactta 60

tgagtcgggtg tcgctggctg tctttggcag catgtgcacg ccttctctat tcatccacac 120

accctgtgc acttgtgaga ctggaggccg taaaaagcct tcagtgctaa attcatatac 180

aaactcattt aactgcactg aatgtacttg atgtaacgca tcattacact 230

<210> 28

<211> 152

<212> DNA

<213> Rhizoctonia sp.


<400> 28

agagtttggt tgtagctggt ccctaattaa cttgggggca tgtgcacacc tttctctttc 60

atcccatata cacctgtgca cctgtgagac agatgttttc tagaggggga aggaacttta 120

ttggacctac tctccttgga ccttctgtct ac 152

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☒ Scanned copy is best available.

*Some drawing  
figures are dark.*

09744503.07301  
TOTET 20 20544260